



Some facts about the Torne and Kalix River Basins

**A contribution to the NEWBALTIC II
workshop in Abisko**

June 1999

Bengt Carlsson

Some facts about the Torne and Kalix River Basins

**A contribution to the NEWBALTIC II
workshop in Abisko**

June 1999

Bengt Carlsson

Some facts about the Torne and Kalix River Basins

A Contribution to the NEWBALTIC II workshop in Abisko, June 1999

An overview

The Torne and Kalix rivers rise in the uppermost north west part of Sweden and their basins are situated between approximately 65°50'N and 69°N as shown in Figure 1. The size basin of the two river basins is about 58 000 km² of which 14 500 km² (25%) is situated in Finland and 350 km² (0.6%) in Norway. The remaining, 43 150 km², makes up 9.7% of the area of Sweden. Data presented in this paper originate from Swedish measurements. Finnish data of snow and evaporation from the east of the Torne River basin can be found in the Finnish Hydrological Yearbook (1994), which also includes some statistics for the period 1961-90.

The density of population is very low, about three people per square kilometre. People are mostly living in the river valleys near the coast, the towns of Haparanda, Torneo and Kalix, or in ore-mining centres like Kiruna.

Most of the basin can be characterised as a flat forest and mire area. The mountains in the north west part compose only 7-8 % of the total drainage basin. The runoff regime is thus characterised by a flow maximum produced by the snowmelt in the forest and swamp region. The mountains are very old. They were formed during the paleozoic era 300 000 000 years ago. As can be seen in Figure 3 at least 1/3 of the basin area is situated below 200masl and also about 1/3 above the 500m elevation line. Topography is further described in Figure 3.

As can be seen from Table 1, the Torne River is number two and the Kalix River number 10 in drainage area order of the Swedish rivers. Due to dominating south-west winds and orographic effects, the Abisko region has a proportionately low precipitation. The specific runoff of the Torne River is thus lower than other Swedish rivers in this part of Sweden.

Table 1. The Torne and Kalix river basins related to the ten greatest Swedish river basins.

Basin	Drainage area (km ²)	Runoff (m ³ /s)	Runoff (mm)	Spec. runoff (l/s*km ²)
The Göta River	50 120	574	360	11.5
The Torne River	40 160	373	310	11
Ångermanälven	31 860	474	490	15.5
Dalälven	28 950	353	380	12.2
Umeälven	26 820	432	520	16
Indalsälven	26 710	437	530	17
The Lule River	25 240	494	610	19.8
Mälaren-Norrström	22 650	162	230	7.2
The Ljusnan River	19 830	228	360	11.5
The Kalix River	18 130	280	500	12.1




Figure 1. The Torne and Kalix rivers and their drainage basin. The map shows the main rivers and the largest tributaries. See also the water divides in Figure 4. (from Hjort 1971)

The Torne River and its tributary Muonio River form the border between Sweden and Finland. The basin area of the Torne River is more than 40 000km². The Torne River drains the largest mountain lake in Sweden, the Torneträsk. Its area is 322 km² and the depth is at most 168m. From Torneträsk to the Bothnian Bay the length of the river is 420 km. The largest tributaries are from north-west to south-east Rautasätno, Vittangi, Lainio, Muonio and Tengeliönjoki rivers.

The source of the Kalix River is the Kebnekaise mountains (2117 masl). The drainage area is some 18 000km². The most important tributaries are Kaitum, Linaälv-Ängesån. The length from source to mouth is about 450km.

Bifurcations are an important element in the drainage system. Through the bifurcation at Junosuando/Tärendö an average of 57% of the discharge is drawn off to the Kalix River. This means 83 m³/s which corresponds to 22% of the total discharge at the mouth. Near the mouth at Karunki the tributary Liankanjoki in average draws off 23% of the discharge. This implies that the mean runoff at Tornio/Haparanda is somewhat less than at Karunki. If the bifurcation at Tärendö is taken into consideration the size of the Torne River Basin becomes about 31 300 km² and the Kalix River Basin about 27 000 km².

It is said that the kind of natural bifurcation at Junosuando/Tärendö is very rare. In fact there is only one more in the world, namely the Casiquiare between the Orinoco and Rio Negro rivers in South America.

Both the Torne and Kalix Rivers should be considered as unregulated. There are some small dams but these do not affect the general runoff. Formerly they were used to regulate the water level for floating, that is, water born transport of timber.




Figure 2. Proposed regulation 1979 of the Torne River.

In the 1970's there was a serious discussion whether the last unregulated four big rivers in Sweden should be developed for hydropower production or not. The Lule and Kalix rivers were two of these rivers. The other two were Pite River and Vindelälven. In 1979 two governmental organisations (Statens Vattenfallsverk and Svenska Kraftverksföreningen) presented their official report which proposed the building of 10 to 15 hydropower stations in each of Torne and Kalix Rivers. This implied a fight between people for and against the hydropower development. As we now know the river environmentalists won this fight. The

development projects were never carried through and we can thus today use these river basins as good examples of unregulated areas in northern Scandinavia.






Figure 3. Topography of the river basins. Two main regions can be distinguished. In the north-west we have a region of high mountains with an altitude of more than 1000 m a s l. In the middle and south-east part there are low mountains, forests and mires. The area of high mountains is less than 10% of the total area and is mainly situated in the upper Kalix River basin. Typical for the basin is the very flat land in the lowermost part and the dales along the river. (from Hjort 1971)




Figure 4. Torne and Kalix River basins with meteorological and stream gauging stations.



Annual total precipitation (mm) 1951-80.



Annual snow precipitation (mm water) 1941-70)



Annual mean temperature (°C) 1951-80.

Figure 5. Annual temperature and precipitation in the Torne River basin. (from Zachrisson 1989)

Near the coast the annual temperature is about $+1^{\circ}\text{C}$. From here it decreases to -3.5°C in the north-west. Around the Torne Träsk it is somewhat milder. The precipitation, which is corrected for windlosses, is mostly around 600-700mm. But also here the area around the Torne Träsk is different as the precipitation is 500mm or less. The gradient towards the highest precipitation, 1600mm, in the mountains in north west is very steep. Days with snow cover increase from about 175 at the coast to 225 in the mountains. In Haparanda the snow has normally melted by the beginning of May.

The maps in Figure 5 are based on the stations in figure 4. See also the time series of temperature and precipitation from the three stations presented in Figure 7.




Figure 6. Specific runoff has been modelled for all Sweden in areas of 25x25km and the values are then connected with isolines. This Figure shows an area covering basins of the Torne and Kalix Rivers.





Figure 7. Annual means and linear fits for precipitation and temperature 1900-1997 in Haparanda (black), Karesuando (red) and Abisko (blue). Are there some trends in these data? Both temperature and precipitation show a small increase in time, but only the rise in precipitation at Karusuando seems to be significant.




Figure 8. Runoff at the station Karunki (Kukkolankoski) 1911-1998, 25 km from the mouth in the Bothnian Bay. Like in the precipitation there may be a weak increasing tendency in the runoff and more in the maximum than in the minimum flows. Notice the ten times greater scale in the minimum values than for the other data. The rise had not been able to be detected with the same scale.





Figure 9. Runoff-modelling of the two rivers. The Torne (top) and Kalix (bottom) Rivers are both set up and modelled with the HBV-model. The HBV-model use 20 sub basins in each of the two main basins. This example from september 1994 to December 1997 shows that the modelling of the recession in the Torne River is inadequate and must be revised.




Figure 10. Daily data of air temperature, precipitation, snow depths and soil temperatures at three depths. Measurements in the Torneträsk area has been carried out since at least 1903, when the Abisko Scientific Research Station was established. Since 1935 it has been under the auspices of The Royal Swedish Academy of Sciences. Some of the long term data sets are presented on the internet, www.ans.kiruna.se, from where this data set, with exception of snow depth (Tjust 1999), is taken.

The sampling site is situated in the mountain birch region not far from the research station. It is obvious that a lot of snow implies a higher soil temperature during the wintertime. The importance of the air temperature at the beginning of snow covered period can be studied for instance by comparing the winters 1991/92 and 1992/93.




Figure 11. Ice thickness in lake Torne Träsk 1960-1990, all measurements about once a week. A great and deep lake situated in a valley between high mountains, mild winters and low precipitation make the ice situation of the lake Torne Träsk somewhat special. The ice is thus laid almost two month later then in other lakes in the basin. The ice thickness is considerable. In the winter 1965/66 the so far highest ice thickness, 110 cm, was registered. Depending on the low precipitation, i.e. not much snow, the ice consists normally mainly of clear ice with little snow ice (Eklund 1998).




Figure 12. Due to the above mentioned climatological and geographical conditions the ice is laid almost two months later in lake Torne Träsk then in the adjacent lakes. Lake: •.




Figure 12. The day of ice break-up in the Torne River at Haparanda has been recorded since 1693. This Figure shows the 10-year mean deviation from the average 1801-1900. Positive values mean earlier ice break-up. Thus the present ice break-up occurs about 8 days earlier than before 1750. A milder climate in the spring is one reason for the change in time of ice break-up.

Acknowledgements

This work was carried out within the NEWBALTIC II project with funds from European Commission's programme Environment and Climate (Contract No ENV4-CT97-0626) and the Swedish Meteorological and Hydrological Institute.

References

Eklund, A., (1998)

Istjocklek på sjöar-en statistisk bearbetning av SMHI's mätningar. (Ice thickness on lakes – a statistical treatment of SMHI's measurements. In Swedish)

SMHI Hydrologi no 76

Hjort, S., (1971)

The Torne and Kalix Rivers. (in Swedish, English summary)

University of Uppsala , Dep. of Physical Geography

UNGI Report no 12

Meteorological and Hydrological data bases at SMHI

Statens Vattenfallsverk och Svenska Kraftverksföreningen. (1979)

Fortsatt Vattenkraftsutbyggnad,

Redovisning till Konsekvensutredningen (in Swedish)

Tjust, M., (1999)

Abisko Scientific Research Station. Personal communication.

Zachrisson, G., (1989)

Svåra islossningar i Torneälven. (Ice break-up in the Torne River, in Swedish)

SMHI Hydrologi nr 25

SMHI ger ut sex rapportserier. Tre av dessa, R-serierna är avsedda för internationell publik och skrivs därför oftast på engelska. I de övriga serierna används det svenska språket.

Seriernas namn	Publiceras sedan
RMK (Rapport Meteorologi och Klimatologi)	1974
RH (Rapport Hydrologi)	1990
RO (Rapport Oceanografi)	1986
METEOROLOGI	1985
HYDROLOGI	1985
OCEANOGRAFI	1985

I serien HYDROLOGI har tidigare utgivits:

- | | |
|--|---|
| <p>1 Bengt Carlsson (1985)
Hydrokemiska data från de svenska fältforskningsområdena.</p> <p>2 Martin Häggström och Magnus Persson (1986)
Utvärdering av 1985 års vårflödesprognoser.</p> <p>3 Sten Bergström, Ulf Ehlin, SMHI, och Per-Eric Ohlsson, VASO (1986)
Riktlinjer och praxis vid dimensionering av utskov och dammar i USA. Rapport från en studieresa i oktober 1985.</p> <p>4 Barbro Johansson, Erland Bergstrand och Torbjörn Jutman (1986)
Skåneprojektet - Hydrologisk och oceanografisk information för vattenplanering - Ett pilotprojekt.</p> <p>5 Martin Häggström (1986)
Översiktlig sammanställning av den geografiska fördelningen av skador främst på dammar i samband med septemberflödet 1985.</p> <p>6 Barbro Johansson (1986)
Vattenföringsberäkningar i Södermanlands län - ett försöksprojekt.</p> <p>7 Maja Brandt (1986)
Areella snöstudier.</p> <p>8 Bengt Carlsson, Sten Bergström, Maja Brandt och Göran Lindström (1987)
PULS-modellen: Struktur och tillämpningar.</p> | <p>9 Lennart Funkquist (1987)
Numerisk beräkning av vågor i kraftverksdammar.</p> <p>10 Barbro Johansson, Magnus Persson, Enrique Aranibar and Robert Llobet (1987)
Application of the HBV model to Bolivian basins.</p> <p>11 Cecilia Ambjörn, Enrique Aranibar and Roberto Llobet (1987)
Monthly streamflow simulation in Bolivian basins with a stochastic model.</p> <p>12 Kurt Ehlert, Torbjörn Lindkvist och Todor Milanov (1987)
De svenska huvudvattendragens namn och mynningspunkter.</p> <p>13 Göran Lindström (1987)
Analys av avrinningsserier för uppskattning av effektivt regn.</p> <p>14 Maja Brandt, Sten Bergström, Marie Gardelin och Göran Lindström (1987)
Modellberäkning av extrem effektiv nederbörd.</p> <p>15 Håkan Danielsson och Torbjörn Lindkvist (1987)
Sjökarte- och sjöuppgifter. Register 1987.</p> <p>16 Martin Häggström och Magnus Persson (1987)
Utvärdering av 1986 års vårflödesprognoser.</p> |
|--|---|

- 17 Bertil Eriksson, Barbro Johansson, Katarina Losjö och Haldo Vedin (1987)
Skogsskador - klimat.
- 18 Maja Brandt (1987)
Bestämning av optimalt klimatstationsnät för hydrologiska prognoser.
- 19 Martin Häggström och Magnus Persson (1988)
Utvärdering av 1987 års vårflödesprognoser.
- 20 Todor Milanov (1988)
Frys förluster av vatten.
- 21 Martin Häggström, Göran Lindström, Luz Amelia Sandoval and Maria Elvira Vega (1988)
Application of the HBV model to the upper Río Cauca basin.
- 22 Mats Moberg och Maja Brandt (1988)
Snökartläggning med satellitdata i Kultsjöns avrinningsområde.
- 23 Martin Gotthardsson och Sten Lindell (1989)
Hydrologiska stationsnät 1989. Svenskt Vattenarkiv.
- 24 Martin Häggström, Göran Lindström, Luz Amelia Sandoval y Maria Elvira Vega (1989)
Aplicacion del modelo HBV a la cuenca superior del Río Cauca.
- 25 Gun Zachrisson (1989)
Svåra islossningar i Torneälven. Förslag till skadeförebyggande åtgärder.
- 26 Martin Häggström (1989)
Anpassning av HBV-modellen till Torneälven.
- 27 Martin Häggström and Göran Lindström (1990)
Application of the HBV model for flood forecasting in six Central American rivers.
- 28 Sten Bergström (1990)
Parametervärden för HBV-modellen i Sverige. Erfarenheter från modellkalibreringar under perioden 1975 - 1989.
- 29 Urban Svensson och Ingemar Holmström (1990)
Spridningsstudier i Glan.
- 30 Torbjörn Jutman (1991)
Analys av avrinningens trender i Sverige.
- 31 Mercedes Rodriguez, Barbro Johansson, Göran Lindström, Eduardo Planos y Alfredo Remont (1991)
Aplicacion del modelo HBV a la cuenca del Río Cauto en Cuba.
- 32 Erik Arnér (1991)
Simulering av vårflöden med HBV-modellen.
- 33 Maja Brandt (1991)
Snömätning med georadar och snötaxeringar i övre Luleälven.
- 34 Bent Göransson, Maja Brandt och Hans Bertil Wittgren (1991)
Markläckage och vattendragstransport av kväve och fosfor i Roxen/Glan-systemet, Östergötland.
- 35 Ulf Ehlin och Per-Eric Ohlsson, VASO (1991)
Utbyggd hydrologisk prognos- och varningstjänst.
Rapport från studieresa i USA
1991-04-22--30.
- 36 Martin Gotthardsson, Pia Rystam och Sven-Erik Westman (1992)
Hydrologiska stationsnät 1992/Hydrological network. Svenskt Vattenarkiv.
- 37 Maja Brandt (1992)
Skogens inverkan på vattenbalansen.
- 38 Joakim Harlin, Göran Lindström, Mikael Sundby (SMHI) och Claes-Olof Brandesten (Vattenfall Hydropower AB) (1992)
Känslighetsanalys av Flödeskommitténs riktlinjer för dimensionering av hel älv.
- 39 Sten Lindell (1993)
Realtidsbestämning av arealnederbörd.
- 40 Svenskt Vattenarkiv (1995)
Vattenföring i Sverige. Del 1. Vattendrag till Bottenviken.
- 41 Svenskt Vattenarkiv (1995)
Vattenföring i Sverige. Del 2. Vattendrag till Bottenhavet.
- 42 Svenskt Vattenarkiv (1993)
Vattenföring i Sverige. Del 3. Vattendrag till Egentliga Östersjön.

- 43 Svenskt Vattenarkiv (1994)
Vattenföring i Sverige. Del 4. Vattendrag till Västerhavet.
- 44 Martin Häggström och Jörgen Sahlberg (1993)
Analys av snösmältningsförlopp.
- 45 Magnus Persson (1993)
Utnyttjande av temperaturens persistens vid beräkning av volymsprognoser med HBV-modellen.
- 46 Göran Lindström, Joakim Harlin och Judith Olofsson (1993)
Uppföljning av Flödeskommitténs riktlinjer.
- 47 Bengt Carlsson (1993)
Alkalinitets- och pH-förändringar i Ume-älven orsakade av minimitappning.
- 48 Håkan Sanner, Joakim Harlin and Magnus Persson (1994)
Application of the HBV model to the Upper Indus River for inflow forecasting to the Tarbela dam.
- 49 Maja Brandt, Torbjörn Jutman och Hans Alexandersson (1994)
Sveriges vattenbalans. Årsmedelvärden 1961 - 1990 av nederbörd, avdunstning och avrinning.
- 50 Svenskt Vattenarkiv (1994)
Avrinningsområden i Sverige. Del 3. Vattendrag till Egentliga Östersjön och Öresund.
- 51 Martin Gotthardsson (1994)
Svenskt Vattenarkiv. Översvämningskänsliga områden i Sverige.
- 52 Åsa Evremer (1994)
Avdunstningens höjdberoende i svenska fjällområden bestämd ur vattenbalans och med modellering.
- 53 Magnus Edström och Pia Rystam (1994)
FFO - Stationsnät för fältforskningsområden 1994.
- 54 Zhang Xingnan (1994)
A comparative study of the HBV model and development of an automatic calibration scheme.
- 55 Svenskt Vattenarkiv (1994)
Svenskt dammregister - Södra Sverige.
- 56 Svenskt Vattenarkiv (1995)
Svenskt dammregister - Norra Sverige.
- 57 Martin Häggström (1994)
Snökartering i svenska fjällområdet med NOAA-satellitbilder.
- 58 Hans Bertil Wittgren (1995)
Kvävetransport till Slätbaken från Söderköpingsåns avrinningsområde
- 59 Ola Pettersson (1995)
Vattenbalans för fältforskningsområden.
- 60 Barbro Johansson, Katarina Losjö, Nils Sjödin, Remigio Chikwanha and Joseph Merka (1995)
Assessment of surface water resources in the Manyame catchment - Zimbabwe.
- 61 Behzad Kouchehi (1995)
Älvtemperaturers variationer i Sverige under en tioårsperiod.
- 62 Svenskt Vattenarkiv (1995)
Sänkta och torrlagda sjöar.
- 63 Malin Kanth (1995)
Hydrokemi i fältforskningsområden.
- 64 Mikael Sundby, Rikard Lidén, Nils Sjödin, Helmer Rodriguez, Enrique Aranibar (1995)
Hydrometeorological Monitoring and Modelling for Water Resources Development and Hydropower Optimisation in Bolivia.
- 65 Maja Brandt, Kurt Ehlert (1996)
Avrinningen från Sverige till omgivande hav.
- 66 Sten Lindell, Håkan Sanner, Irena Nikolushkina, Inita Stikute (1996)
Application of the integrated hydrological modelling system IHMS-HBV to pilot basin in Latvia
- 67 Sten Lindell, Bengt Carlsson, Håkan Sanner, Alvina Reihan, Rimma Vedom (1996)
Application of the integrated hydrological modelling system IHMS-HBV to pilot basin in Estonia
- 68 Sara Larsson, Rikard Lidén (1996)
Stationstäthet och hydrologiska prognoser.

- 69 Maja Brandt (1996)
Sedimenttransport i svenska vattendrag
exempel från 1967-1994.
- 70 Svenskt Vattenarkiv (1996)
Avrinningsområden i Sverige. Del 4.
Vattendrag till Västerhavet.
- 71 Svenskt Vattenarkiv (1996)
Svenskt sjöregister. 2 delar
- 72 Sten Lindell, Lars O Ericsson, Håkan Sanner,
Karin Göransson SMHI
Malgorzata Mierkiewicz , Andrzej
Kadlubowski, IMGW (1997)
Integrated Hydrological Monitoring and
Forecasting System for the Vistula River
Basin. Final report.
- 73 Maja Brandt, Gun Grahn (1998)
Avdunstning och avrinningskoefficient i
Sverige 1961-1990. Beräkningar med HBV-
modellen.
- 74 Anna Eklund (1998)
Vattentemperaturer i sjöar, sommar och vinter
- resultat från SMHIs mätningar.
- 75 Barbro Johansson, Magnus Edström, Katarina
Losjö och Sten Bergström (1998)
Analys och beräkning av
snösmältningsförlopp.
- 76 Anna Eklund (1998)
Istjocklek på sjöar.
- 77 Björn Bringfelt (1998)
An evapotranspiration model using SYNOP
weather observations in the Penman-Monteith
equation
- 78 Svenskt Vattenarkiv (1998)
Avrinningsområden i Sverige. Del 2
Vattendrag till Bottenhavet.
- 79 Maja Brandt, Anna Eklund (1999)
Snöns vatteninnehåll Modellberäkningar
och statistik för Sverige



Swedish Meteorological and Hydrological Institute
SE 601 76 Norrköping, Sweden.
Tel +46 11-495 80 00. Fax +46 11-495 80 01