

PAPER • OPEN ACCESS

The effect of current and future maternal exposure to near-surface ozone on preterm birth in 30 European countries—an EU-wide health impact assessment

To cite this article: Johan Ekland *et al* 2021 *Environ. Res. Lett.* **16** 055005

View the [article online](#) for updates and enhancements.

ENVIRONMENTAL RESEARCH
LETTERS

LETTER

OPEN ACCESS

RECEIVED
18 August 2020REVISED
8 February 2021ACCEPTED FOR PUBLICATION
16 February 2021PUBLISHED
20 April 2021

Original content from
this work may be used
under the terms of the
[Creative Commons
Attribution 4.0 licence](#).

Any further distribution
of this work must
maintain attribution to
the author(s) and the title
of the work, journal
citation and DOI.



The effect of current and future maternal exposure to near-surface ozone on preterm birth in 30 European countries—an EU-wide health impact assessment

Johan Ekland¹, David Olsson², Bertil Forsberg², Camilla Andersson³ and Hans Orru^{2,4} ¹ Women's and Children's Health, Uppsala University, Sweden, Uppsala² Public Health and Clinical Medicine, Umea University, Umea, Sweden³ Research department, Swedish Meteorological and Hydrological Institute, Norrköping, Sweden⁴ Family Medicine and Public Health, University of Tartu, Tartu, EstoniaE-mail: Hans.Orru@umu.se**Keywords:** air pollution, preterm birth, emissions, climate change, childrenSupplementary material for this article is available [online](#)

Abstract

Preterm birth is the largest contributor to neonatal mortality globally and it is also associated with several adverse health outcomes. Recent studies have found an association between maternal exposure to air pollution and an increased risk for preterm birth. As a constituent of air pollution, ozone is a highly reactive molecule with several negative health effects when present near earth's surface. This health impact assessment aims to estimate the proportion of preterm births—in current and future situations—attributable to maternal ozone exposure in 30 European countries (EU30). A literature search was performed using relevant keywords, followed by meta-analysis with STATA software in which five studies investigating exposure-response relationship of interest were included. The attributable proportion, and number of cases, was modelled with the software AirQ+ against current and future European ozone concentrations. According to our meta-analysis, the relative risk for giving birth preterm was calculated to 1.027 (95% CI 1.009–1.046) per $10 \mu\text{g m}^{-3}$ increase in ozone concentration. This rendered 7.1% (95% CI 2.5–11.7) of preterm births attributable to maternal ozone exposure to in EU30 during 2010, which is equal to approximately 27 900 cases. By 2050, the projected decrease in ozone precursor emissions rendered an estimated 30% decrease of ozone attributable preterm births. Not taking emission change into account, due to climate change the ozone-related preterm birth burden might slightly increase by 2050 in Central and Southern Europe, and decrease in Eastern and Northern Europe. In summation, these numbers make a substantial impact on public health.

1. Introduction

Preterm birth is commonly defined as birth before the gestational age of 37 weeks, and predisposes children several hazards with both immediate and future health. In 2015 alone, preterm birth caused globally approximately 1 million deaths before the age of 5 [1]. This equates to more than 2700 premature deaths daily, ranking complications emanating from preterm birth the largest contributor of neonatal death globally [2]; however, it is not only the alarming number of mortalities caused by preterm birth which pose a global threat to children's health. The infants that do survive will face an increased risk of

a wide range of etiologies including cognitive, neurological, immunological, and gastrointestinal dysfunctions, along with chronic morbidities such as asthma and type two diabetes later in life [3–5]. Therefore, it is important to acknowledge that the hazardous consequences of prematurity are not necessarily limited to a defined time-period surrounding childhood, but may likewise affect an individual's health and life far into adult years [6].

Several risk factors surrounding preterm birth have previously been established, including facets of basic maternal lifestyle choices such as physical activity, diet, alcohol consumption, and tobacco use. The role of more complex contributors, such as genetics

and environmental factors, is less explored, albeit some associations have been suggested [3].

1.1. Air pollution and near-surface ozone

The World Health Organization (WHO) currently describes air pollution as the biggest environmental risk to health as the proportion of worlds' population living in areas where air pollutants exceed the safe levels, is estimated to 92% [7]. Several studies have recognized that air pollution has adverse effects on health by showing associations between air pollution exposure and morbidity, foremost of cardiovascular and respiratory etiology [8]. Air pollution—as mixture of various substances—include near-surface, sometimes called ground-level or tropospheric, ozone. Unlike several other air pollutants, near-surface ozone is not directly emitted, but rather photochemically formed in complex reactions when nitrogen oxide (NO_x) is exposed to sunlight in the presence of volatile organic compounds (VOCs) [9]. Hence, apart from the levels of precursor elements such as NO_x and VOCs, the rate of near-surface ozone formation is dependent on meteorological conditions such as the amount of sunlight, wind, and, most importantly, temperature on the surface [10]. As understood by a panoply of scientists, we will experience higher global temperatures with increased health effects in the future due to anthropogenic climate change [11]. As described by Doherty *et al* [12], many of the meteorological consequences of climate change, such as higher temperature, less cloud coverage, and less rainfall can synergistically contribute to an increase in near-surface ozone concentration, due to more efficient ozone formation and increases in natural emissions of VOCs. On the contrary, it should be noted that the emission of anthropogenic ozone precursors have decreased in all EU-member countries and are projected to decrease further in the future [13, 14].

In 2005, the EU has set a target value and WHO a guideline for the ozone concentration in ambient air for a daily 8 h maximum of 120 and 100 micrograms per cubic meter ($\mu\text{g m}^{-3}$), respectively [15]. However, a growing number of studies are finding adverse health effects from ozone exposures at even lower concentrations [16–20]. For instance, a study from the US by Bell *et al* [17] found associations between premature mortality and ozone exposures near typical background ozone levels at $20 \mu\text{g m}^{-3}$, thus suggesting that there are almost no safe levels of near-surface ozone. For this reason, recommendations are starting to shift in regard to what cut-off value to use when assessing the health impacts of ozone. Amann *et al* [16] estimated earlier approximately 21 000 premature deaths were associated with ozone concentrations above $70 \mu\text{g m}^{-3}$ in 25 European countries; however, with SOMO25 and inclusion of long-term effects, this estimation could rise to 55 000 [21]. As reviewed by Orru *et al* [22], different studies have

applied different cut-off values and to reduce these divergencies, we have in this study chosen to adopt an ozone exposure cut-off at $50 \mu\text{g m}^{-3}$ expressed as sum of means over 25 ppb (SOMO25). It is the yearly sum of the daily maximum of 8 h running average over 25 ppb.

A majority of the studies tackling the health effects of air pollution have traditionally focused on general mortality, or respiratory and vascular morbidity [23–25]. However, alternative endpoints such as cognitive development and reproductive health, have also been suggested [8]. Several studies have shown associations between ozone exposure, specifically, and negative birth outcomes [26–28]. The theories regarding the pathogenesis of ozone exposure are related to an ozone molecule's ability to promote free radical reactions in the body [29]. As free radicals come in contact with target molecules, they may become inactivated, initiating cellular damage through oxidation progressions referred to as oxidative stress [29]. It is suggested that oxidative stress, along with hemodynamics disruptions and inflammation in the mother, impairs the delivery of oxygen and nutrients to the foetus, thus causing a disturbed gestational process [30]. The increased levels of proinflammatory mediators resulting from oxidative stress can subsequently provoke, or increase, the risk for giving birth preterm [30–32]. This theory is supported by additional literature showing that systemic inflammation can trigger a preterm delivery [33]. A biological marker of inflammation in the body is the plasma level of C-reactive protein (CRP). In early pregnancy, exposure to certain air pollutants, ozone included, has been shown to increase CRP levels and then act as a potential trigger for preterm birth [34].

The existing knowledge regarding the extent of ozone related birth effects' is limited thus, to the best of our knowledge, this is the first continent-wide health impact assessment (HIA) quantifying maternal ozone exposure and the number of attributable cases of preterm birth. It aims to estimate the attributable proportion (AP) of preterm births that occur due to maternal ozone exposure during the first trimester of pregnancy. Moreover, we aim to quantify how ozone concentration due to changes in climate and emission levels change, and we estimate the AP for year 2050 in 30 European countries (EU30).

2. Materials and methods

2.1. HIA

A HIA utilizes a methodology for quantifying the health outcomes of a certain exposure [35]. The procedure follows a series of steps in which, firstly, an exposure-response function is selected—usually expressed as a relative risk. This exposure-response function is then applied to an outcome baseline to estimate the AP and number of cases caused by a certain exposure, e.g. near-surface ozone [36].

This methodology has been utilized by several other environmental health studies [37–39] and also by a similar study which investigated preterm birth as an outcome, but particulate matter as an exposure constituent [40].

2.2. Near-surface ozone modelling

First, to determine human exposure to near-surface ozone in Europe, the multi-scale atmospheric transport and chemistry (MATCH) model [41–43], that is Eulerian chemistry transport model (CTM) was applied at a 50×50 km horizontal resolution. The model used here includes ozone and particle forming chemistry considering ca 60 chemical species [44]. Meteorological 3D information is updated every hour in the model with an impact on concentrations in thermal- and photochemical reactions (e.g. temperature, cloud cover, solar radiation, humidity), transport, deposition and boundary layer processes (e.g. wind, temperature profile, precipitation, humidity, solar radiation), and the natural emissions of isoprene calculated online (e.g. temperature, solar radiation) following the Simpson *et al* [45] E-94 methodology, while terpene emissions are not treated in this model version. Isoprene chemistry is based on an adapted version of the Carter one-product mechanism [44, 46], while other gas-phase chemistry is based on the EMEP MSC-W EMChem09 scheme of Simpson *et al* [47] with reaction rate updates following the recommendations of International Union of Pure and Applied Chemistry (IUPAC) [41, 48]. The photolysis rates depend on photolytically active radiation, which is calculated from latitude, the time of day, and cloud cover. The dry deposition is modelled through a resistance approach. The dry deposition to stomata—modelled as described by Andersson and Engardt [49]—is dependent on soil moisture, temperature, solar radiation, and the humidity of the air. Thus, many processes in the model are impacted by climate change. MATCH is widely used for various air quality and, specifically, ozone research and operational services, such as daily European operational air quality forecasting [50] and national Swedish environmental surveillance (e.g. [51]) of ozone and deposition. The ozone performance of MATCH is commonly among the best in comparison with other European CTMs (e.g. [52]). The specific set up and evaluation of ozone were described in detail previously [13, 14, 21, 53, 54], but briefly recaptured here. Ozone concentrations from this set up were evaluated for the present situation compared to airbase measurement sites (see [54]), where MATCH had the lowest root mean square error of the four CTMs and a mean annual and summer bias of 1 ppb(v) and -1.1 ppb(v), respectively. MATCH was forced by the global climate model EC-EARTH and downscaled with the Rossby Climate regional climate model RCA4 [55]. They were both forced by the representative concentration pathway scenario RCP4.5 [56]. MATCH was

also forced on the boundaries by the LMDz-INCA global model [57] and by anthropogenic air pollution precursor emissions of ECLIPSE v4a for the present (years 2005–2010) and the future (year 2050, current legislation). The evolution is considered coherent with the future evolution of the RCP4.5 scenario at 2050 [14].

In the current study the SOMO25 (the highest daily 8 h average exceeding 25 parts per billion (ppb) summarized from day 1 to 365) values have been applied. The population-weighted exposure (E_w) has been calculated by multiplying the SOMO25 concentration within each 50×50 km grid (E_i) by the number of individuals residing within the same grid (POP_i). The sum of the concentration of all grids was then divided by the total population of respective country (POP_{tot}) as: $E_w = \frac{\sum(E_i \times POP_i)}{POP_{tot}}$.

2.3. Preterm birth data



Secondly, the national baseline incidence rate of preterm birth for year 2010 in EU30 was obtained from the United Nations Statistics Division database [58], where data has been collected through examination of national and local administrative reporting registers, civil registration forms, and surveys. Data on current (2010) population size for each country in EU30 was likewise extracted from the WHO database [59] and for future (2050) populations Eurostat population projections [60].

2.4. Exposure-response function

To determine a pooled exposure-response function, a systematic literature review was conducted (figure 1). The scientific databases PubMed, CINAHL and Cochrane library were utilized in October 2017 using the terms ‘Ozone’ AND ‘Preterm birth,’ ‘Ozone’ AND ‘Preterm delivery,’ ‘Air pollution’ AND ‘Premature birth.’ Publications which met the inclusion criteria were exposure being ozone and outcome being preterm birth, which were screened through an abstract and full text review. Between the 14 examined studies [61–74], significant divergencies in exposure variables were observed. The most fundamental differences were differences in the exposure period, either first trimester, third trimester or whole pregnancy, and the type of ozone exposure value (1 h, 8 h and 24 h maximum). Therefore, a second set of inclusion criteria was employed. Studies assessing first trimester ozone exposure were measured at an 8 h maximum due to the larger number of studied phenomena and more profound effects with either first or third trimester exposure data, thus leaving five studies [63, 64, 69, 71, 74] (figures 1 and 2).

2.5. Meta-analysis

The meta-analysis of the five studies was done in the statistical software environment of STATA 13 using the ‘metan’ package [75]. The software designated a



certain impact on the overall effect size depending on the level of uncertainty in each study. This meant that the wider confidence interval a certain study carried, the less influence it would have over the overall effect size (figure 2). All studies, which did not present an effect size per $10 \mu\text{g m}^{-3}$ increase in ozone concentration, were recalculated accordingly. The meta-analysis effect model used was the DerSimonian and Laird mixed-effect (random effect) model [76, 77] (supplementary material (available online at stacks.iop.org/ERL/16/055005/mmedia)).

2.6. Calculation of attributable effects

To estimate the proportion and number of preterm births caused by maternal ozone exposure in Europe, the exposure-response function was applied to the ozone levels in each respective country. The AP and number of cases were calculated using the WHO developed software *AirQ+*. This WHO developed software can be used to quantify health effects of certain air pollutants, including ozone. The estimated impact is grounded on a log-linear relationship between relative risk (RR) and concentration levels.

The AP is then estimated as defined: $\text{AP} = \frac{\sum [\text{RR}(c) - 1] \times p(c)}{\sum [\text{RR}(c) \times p(c)]}$,


where $p(c)$ is the proportion of the population.

To generate an estimate of the excess number of cases of preterm birth attributable to ozone exposure (ΔY), the software uses the baseline preterm birth rate (Y_0) and the population at risk (pop), defined as: $\Delta Y = (Y_0 \times \text{pop}) \times (\exp^{\beta \times \Delta X})$, where β represents the effect size and ΔX is the variation in concentration.

3. Results

3.1. Ozone levels

The current population average SOMO25 level for EU30 was $10\,172 \mu\text{g m}^{-3}\cdot\text{d}$, ranging from 6667 to $18\,230 \mu\text{g m}^{-3}\cdot\text{d}$ (figure 3). The highest SOMO25 values were recorded in the Mediterranean region with Malta, Cyprus, Greece, Italy, and Portugal being the five highest exposed countries with a combined SOMO25 average value of $15\,351 \mu\text{g m}^{-3}\cdot\text{d}$, that would mean around $92 \mu\text{g m}^{-3}$ as daily 8 h max mean. The lowest levels were found in the United Kingdom, Finland, and the Baltic States due to low levels during winter period, around $50 \mu\text{g m}^{-3}$ as daily 8 h max mean. The average SOMO25 level for the least exposed countries was $6847 \mu\text{g m}^{-3}\cdot\text{d}$.



In 2050, the overall ozone concentrations based on climate change were projected to be left relatively unchanged when compared to the current EU30 average (an average increase with 0.3%). However, some individual countries saw more tangible climate effects in the form of both increases and decreases. Changes in SOMO25 values due to an altered climate yielded an increase in Central-Europe (e.g. in Luxembourg 12% increase in SOMO25 levels) to a 6%–7% decrease in the Baltic States (Estonia, Latvia and Lithuania).

In the future emissions-based model, all EU30 countries presented significant reductions which in all cases were greater than reductions driven by climate change. The MATCH estimation for future SOMO25 levels based on changes in ozone precursor emissions yielded an EU30 average reduction of 29%, ranging from a SOMO25 decrease by 14% for Cyprus to a 38% reduction for Hungary.

3.2. Pooled ozone effect size estimates from meta-analysis

The studies used for the modelling of an exposure-response function [63, 64, 69, 71, 74] had a sample size range from approximately 28 000 to 1.5 million subjects. The mean daily 8 h maximum ozone concentration, as short-term exposure, varied in studies between 53.1 and 84.5 $\mu\text{g m}^{-3}$. The studies adjusted for a varying set of possible confounders such as age, smoking, parity, co-morbidity, the season of conception, and co-pollutants effects. All studies used similar methodologies for pollutant measurements and statistical analysis, and all used the same maternal exposure window, which was average exposure during the first trimester. The increased risk for giving birth preterm based on ozone exposure were expressed as odds ratios and ranged from 1.01 to 1.18 (95% CI 1.00–1.30) per 10 $\mu\text{g m}^{-3}$ increase in ozone concentration. When weighted and modelled, the respective effect sizes generated a pooled effect size of 1.027 (95% CI 1.009–1.046) per 10 $\mu\text{g m}^{-3}$

increase in the mean daily 8 h maximum ozone concentration (figure 2).

3.3. Number of current preterm births due to near-surface ozone exposure in 2010

The average population proportion attributable to preterm birth caused by ozone was estimated to 7.2% (95% CI 2.5–11.8) (table 1). The largest attributable proportions were found in countries with higher SOMO25 values, i.e. the Mediterranean region where Malta, Cyprus, and Greece had an estimated AP of 12.5% (4.4–20.2), 10.7% (3.7–17.4), and 10.4% (3.6–17.0), respectively. The smallest AP was found in the northern parts of Europe, where SOMO25 values were the lowest. The United Kingdom, Finland, and Lithuania showed an estimated AP of 4.7% (1.6–7.7), 5.1% (1.7–8.4) and 5.2% (1.8–8.7), respectively.

In absolute numbers, more preterm births attributable to ozone were found in countries with larger populations, e.g. France reported the highest total number of attributable preterm births (3889, 95% CI 1353–6340) related to maternal ozone exposure. Conversely, due to its small population, Iceland, who also experienced a relatively small attributable proportion, presented the smallest number of attributable cases: 20 (95% CI 7–34) preterm births attributable to maternal ozone exposure. The total number of attributable cases in EU30 in 2010 was estimated to be 27 871 (95% CI 9611–45 854) premature births.

3.4. Future estimates of preterm births due to near-surface ozone exposure and populations change in 2050

Climate change will only slightly alter the attributable impact, which relates to the moderate changes in SOMO25 values in the same model. If population baseline rates remain unchanged, we expect altogether 424 more cases of preterm birth in EU30. In this analysis, the EU30 average near-surface ozone induced AP of preterm birth is expected to decrease

Table 1. EU30 attributable proportion (AP, %), attributable number of cases (N) with 95% confidence interval (95% CI) due to near-surface ozone exposure (under SOMO25 values) currently (2005–2010) and in the future (2050), taking climate and emissions change into account.

Country	Current			Climate change			Emissions change			Future			Changes combined		
	AP% (95% CI)	N (95% CI)		AP% (95% CI)	N (95% CI)		AP% (95% CI)	N (95% CI)		AP% (95% CI)	N (95% CI)		AP% (95% CI)	N (95% CI)	
Austria	8.5 (2.9–14.0)	738 (255–1209)		8.5 (2.9–13.9)	733 (254–1200)		5.8 (2.0–9.7)	507 (174–838)		8.5 (2.9–14.0)	799 (276–1309)		5.8 (2–9.6)	545 (188–901)	
Belgium	4.6 (1.6–7.7)	470 (161–781)		5.1 (1.7–8.4)	514 (176–852)		3.8 (1.3–6.4)	390 (133–650)		4.6 (1.6–7.7)	531 (182–882)		4.2 (1.4–7)	482 (164–801)	
Bulgaria	9.3 (3.2–15.1)	560 (194–915)		9.1 (3.2–14.9)	551 (191–901)		6.0 (2.0–9.9)	360 (124–595)		9.3 (3.2–15.1)	645 (223–1054)		5.9 (2–9.8)	408 (141–675)	
Switzerland	9.3 (3.2–15.2)	520 (181–850)		9.3 (3.2–15.2)	521 (181–852)		6.5 (2.2–10.8)	366 (126–604)		9.3 (3.2–15.2)	672 (234–1099)		6.5 (2.2–10.8)	474 (163–783)	
Czech Republic	6.7 (2.3–11.1)	550 (189–907)		6.8 (2.3–11.1)	554 (191–914)		4.6 (1.6–7.6)	375 (128–623)		6.7 (2.3–11.1)	556 (191–917)		4.7 (1.6–7.6)	382 (131–635)	
Cyprus	10.7 (3.7–17.4)	157 (55–256)		10.5 (3.7–17.1)	154 (54–251)		9.1 (3.2–14.9)	134 (47–219)		10.7 (3.7–17.4)	127 (45–207)		8.9 (3.2–14.6)	107 (37–174)	
Germany	5.7 (2.0–9.5)	3518 (1206–5821)		6.0 (2.1–10.0)	3690 (1267–6099)		4.2 (1.4–7.0)	2586 (882–4301)		5.7 (2.0–9.5)	3401 (1166–5627)		4.4 (1.5–7.4)	2622 (896–4356)	
Denmark	5.6 (1.9–9.3)	249 (85–412)		5.5 (1.9–9.2)	245 (84–406)		4.1 (1.4–6.8)	182 (62–302)		5.6 (1.9–9.3)	283 (97–468)		4 (1.4–6.7)	203 (70–338)	
Estonia	5.5 (1.9–9.1)	52 (17–86)		5.2 (1.8–8.6)	49 (17–82)		3.8 (1.3–6.3)	36 (12–59)		5.5 (1.9–9.1)	45 (15–75)		3.6 (1.2–0.1)	29 (10–49)	
Spain	8.9 (3.1–14.5)	2988 (1036–4888)		9.0 (3.1–14.8)	3034 (1053–4961)		6.2 (2.1–10.2)	2079 (714–3435)		8.9 (3.1–14.5)	2952 (1023–4829)		6.3 (2.1–10.4)	2086 (717–3444)	
Finland	5.1 (1.7–8.4)	185 (63–306)		4.8 (1.6–7.9)	174 (60–289)		3.6 (1.2–6.0)	131 (44–218)		5.1 (1.7–8.4)	189 (64–313)		3.4 (1.1–5.6)	126 (43–211)	
France	6.8 (2.4–11.3)	3817 (1314–6291)		7.2 (2.5–11.8)	4001 (1379–6586)		4.7 (1.6–7.8)	2605 (890–4326)		6.8 (2.4–11.3)	4159 (1432–6855)		5 (1.7–8.1)	2975 (1018–4935)	
Greece	10.4 (3.6–17.0)	803 (280–1306)		10.4 (3.6–16.8)	797 (278–1297)		7.2 (2.5–11.8)	553 (191–910)		10.4 (3.6–17.0)	648 (226–1055)		7.2 (2.5–11.7)	443 (153–730)	
Hungary	7.2 (2.5–11.8)	583 (201–959)		7.1 (2.4–11.6)	574 (198–946)		4.4 (1.5–7.4)	361 (123–600)		7.2 (2.5–11.8)	496 (171–816)		4.3 (1.4–7.3)	303 (103–504)	
Ireland	5.6 (1.9–9.2)	247 (85–409)		5.5 (1.9–9.1)	245 (84–405)		4.2 (1.4–7.0)	185 (36–308)		5.6 (1.9–9.2)	296 (102–490)		4.1 (1.4–6.9)	220 (43–365)	
Iceland	6.0 (2.0–9.8)	20 (7–34)		5.7 (2.0–9.4)	20 (7–32)		4.6 (1.5–7.7)	16 (5–26)		6.0 (2.0–9.8)	23 (8–39)		4.4 (1.5–7.4)	18 (6–28)	
Italy	9.9 (3.4–16.1)	3889 (1353–6340)		9.9 (3.5–16.2)	3905 (1359–6365)		6.8 (2.3–11.2)	2673 (920–4406)		9.9 (3.4–16.1)	3527 (1227–5750)		6.8 (2.4–11.3)	2434 (838–4012)	
Lithuania	5.2 (1.8–8.7)	112 (38–185)		4.9 (1.7–8.1)	104 (36–173)		3.5 (1.2–5.8)	74 (25–124)		5.2 (1.8–8.7)	73 (25–121)		3.3 (1.1–5.4)	45 (16–76)	
Luxembourg	4.9 (1.7–8.3)	23 (8–38)		5.5 (1.9–9.1)	26 (9–43)		4.2 (1.4–7.0)	20 (7–33)		4.9 (1.7–8.3)	36 (13–60)		4.7 (1.6–7.7)	36 (12–59)	
Latvia	5.2 (1.8–8.7)	50 (17–83)		4.9 (1.7–8.1)	47 (16–78)		3.5 (1.2–5.8)	33 (11–56)		5.2 (1.8–8.7)	33 (11–55)		3.3 (1.1–5.4)	21 (7–35)	
Malta	12.5 (4.4–20.2)	30 (11–49)		12.5 (4.4–20.1)	30 (11–48)		9.0 (3.1–14.8)	22 (8–35)		12.5 (4.4–20.2)	32 (12–52)		9 (3.1–6.7)	23 (8–36)	
Netherlands	5.0 (1.7–8.3)	733 (251–1216)		5.3 (1.8–8.8)	779 (267–1291)		4.3 (1.5–7.1)	628 (214–1045)		5.0 (1.7–8.3)	752 (257–1247)		4.6 (1.6–7.2)	684 (233–1138)	
Norway	6.0 (2.10.0)	222 (76–367)		5.8 (2.0–9.5)	212 (73–351)		4.5 (1.5–7.5)	167 (57–277)		6.0 (2.10.0)	302 (103–499)		4.4 (1.4–6.6)	217 (74–360)	
Poland	5.8 (2.0–9.7)	1689 (579–2793)		5.6 (1.9–9.3)	1630 (559–2699)		3.9 (1.3–6.5)	1130 (385–1881)		5.8 (2.0–9.7)	1474 (505–2438)		3.8 (1.2–6.2)	952 (324–1587)	
Portugal	9.5 (3.3–15.4)	767 (266–1251)		9.8 (3.4–15.9)	790 (275–1289)		6.4 (2.2–10.6)	519 (179–857)		9.5 (3.3–15.4)	687 (238–1120)		6.6 (2.3–10.9)	479 (166–791)	
Romania	8.0 (2.8–13.1)	1186 (410–1947)		7.8 (2.7–12.8)	1162 (401–1909)		5.1 (1.7–8.4)	755 (258–1252)		8.0 (2.8–13.1)	910 (315–1494)		5 (1.6–8.2)	568 (194–942)	
Sweden	5.6 (1.9–9.3)	388 (133–643)		5.4 (1.8–8.9)	373 (128–618)		4.0 (1.4–6.7)	279 (95–464)		5.6 (1.9–9.3)	479 (164–794)		3.9 (1.3–6.4)	331 (113–551)	
Slovenia	9.1 (3.2–14.9)	158 (55–259)		9.1 (3.2–14.9)	158 (55–258)		6.2 (2.1–10.2)	107 (37–176)		9.1 (3.2–14.9)	152 (53–250)		6.2 (2.1–10.2)	103 (36–169)	
Slovakia	7.3 (2.5–12.1)	266 (92–438)		7.2 (2.5–11.9)	261 (90–430)		4.8 (1.6–7.9)	173 (59–288)		7.3 (2.5–12.1)	244 (84–401)		4.7 (1.6–7.8)	155 (53–259)	
United Kingdom	4.7 (1.6–7.7)	2901 (991–4818)		4.7 (1.6–7.9)	2959 (1011–4914)		3.9 (1.3–6.5)	2432 (829–4049)		4.7 (1.6–7.7)	3477 (1188–5774)		3.9 (1.3–6.7)	2973 (1014–4949)	
EU-30	7.2 (2.5–11.8)	27 871 (9611–45 854)		7.1 (2.5–11.7)	28 295 (9761–46 538)		5.1 (1.7–8.4)	19 876 (6801–32 960)		7.2 (2.5–11.8)	28 078 (9681–46 198)		5 (1.7–8.3)	20 328 (6936–33 696)	

from 7.1% to 5.1%, as a consequence of reduced emissions. Subsequently, as the EU population is expected to increase, the effects on preterm birth could increase in total by 1%, whereas the effects decrease a third in Latvia and Lithuania and increase by more than 50% in Luxembourg. Seen over the whole EU30 region and all three factors together, the total number of attributable preterm births in 2050 is expected to be 7543 fewer than in 2010 (table 1).

4. Discussion

This multinational study evaluates the attributable effects of maternal near-surface ozone exposure on preterm birth. From the meta-analysis—which included five studies concerning maternal ozone exposure during the first trimester and the subsequent risk for preterm birth—we estimate the increased risk for preterm birth to be 2.7% per $10 \mu\text{g m}^{-3}$ increase in near-surface ozone concentration. This generated a current average AP of 7.2%, or approximately 27 900 preterm births caused by maternal ozone exposure. Should future ozone concentration adhere to the air pollution emissions, climate scenarios, and population projections used here, nearly 27% fewer preterm births are to be expected in 2050. It is, however, important to acknowledge that the potential effect on the European preterm birth incidence rate is only due to maternal ozone exposure. There are many factors that impact the preterm birth incidence rate in a chosen setting, but none other than maternal ozone exposure is investigated here. Each of these results will be discussed and elaborated upon below.

Firstly, the result from the meta-analysis (the increased RR for preterm birth 1.027 per $10 \mu\text{g m}^{-3}$ increase in ozone exposure during first trimester) can be compared with corresponding studies, where other ozone exposure windows have been used. These studies have yielded RRs between 1.02 and 1.05 [66, 67, 72], which is similar to our results. The elevated risk for preterm birth may also be compared with the risks of exposures to $\text{PM}_{2.5}$. In a meta-analysis comprised from 13 studies, Sun *et al* [78] estimated the elevated risk for preterm birth caused for every $10 \mu\text{g m}^{-3}$ increase in $\text{PM}_{2.5}$ concentration during the entire pregnancy to be 13%; however, extensive heterogeneity between the included studies rendered a rather high level of uncertainty in the study. They applied exposure-response function from Sun *et al* [78] meta-analysis that included only studies fully adjusted for other covariates and pollutants.

Although a majority of studies have found positive associations between air pollution and preterm birth, some studies have found no or negative associations [79, 80]. Likewise, some studies have not been able to establish significant associations between maternal ozone exposure, specifically, and the risk for preterm birth [62, 70, 73]. Stieb *et al*

[80] have discussed that part of this inconsistency may be explained by the heterogenic methodologies between studies as both outcomes and pollutants varied widely, and often multi-pollutant models were utilized. Another complicating factor is that in almost half of the cases of preterm birth, the underlying cause remains unidentified [81], making it reasonable to assume that the driving force behind preterm birth mostly will occur as a result of several interacting factors with varying impact capacity [2]. Moreover, the pathophysiology on how ozone affects the gestational environment surrounding the foetus is not yet fully understood. The mechanism by which preterm birth could be triggered by air pollutants is believed to be related to oxidative stress and inflammation, which in turn would disrupt the delicate processes surrounding gestation [32]. Still, a growing body of scholarship currently focuses on establishing associations for exposure to critical air pollutants in general [1, 82–85] and for ozone exposure in particular [26, 61, 67, 70, 72].

Secondly, we have applied the SOMO25 concept (yearly sum of the daily maximum of 8 h running average over 25 ppb) in our HIA, whereas epidemiological studies have applied actual concentrations (ppb, $\mu\text{g m}^{-3}$). The rationale behind this is that SOMO is a summary estimate—it helps to reduce the uncertainties of the concentration—response function at very low concentrations as its linearity does not extend down to zero and it reflects seasonal cycle, e.g. very low levels at winter and without SOMO we would underestimate the effects [8, 86, 87]. When we accounted for impacts from a cut-off of $50 \mu\text{g m}^{-3}$ (SOMO25), the excess risk caused by maternal near-surface ozone exposure yielded an estimated average 7.2% AP of preterm birth in EU30 in 2010. These results can be interpreted as out of 100 preterm births, seven are (in combination with other risk factors) the consequence of a maternal exposure to ozone during the first trimester of pregnancy. Accumulated across the studied European countries, the attributable number of cases due to near-surface ozone is estimated currently to be approximately 27 900 preterm births. This can be compared to a similar study conducted by Malley *et al* [40] that investigated the global AP of preterm birth caused during the entire pregnancy $\text{PM}_{2.5}$ exposure over $10 \mu\text{g m}^{-3}$. In this study, it was estimated that in Europe, 34 800 preterm births were due to fine particles exposure. The applied exposure-response function was based on a meta-analysis that included only studies fully adjusted for other covariates and pollutants [78], thus we cannot expect overlapping effects with ozone.

Our HIA has applied ozone concentrations exceeding $\sim 50 \mu\text{g m}^{-3}$ (SOMO25) when estimating the impact of ozone on preterm birth. This cut-off is, to some degree, lower compared to previous HIAs which adopted concentrations over $\sim 70 \mu\text{g m}^{-3}$ (SOMO35) [16, 88, 89]. The rationale behind using

SOMO25 as a concentration cut-off is based on the earlier mentioned indications that suggest health effects occur in low ozone concentrations, such as typical background values [17–20]. A similar cut-off was also applied in a recent EU-wide ozone-related mortality impact assessment [21]. Also, in three of the five studies included into our meta-analysis, the ozone levels were close to SOMO25 levels [64, 71, 74]. If the $\sim 70 \mu\text{g m}^{-3}$ (SOMO35) cut-off would have been used, the effect-estimate would be more than two times lower (supplementary material). The difference is especially large if we compare the ‘future emissions’ scenarios as the levels are expected to be mostly below $70 \mu\text{g m}^{-3}$. With SOMO35, only a little more than 4000 preterm births are expected with future emissions. This would probably underestimate the real effects based on current susceptibility [17–20].

Another variable in exposure variation is the estimation of averaging time of exposure, as 1 h or 8 h max or 24 h average. Since ground-level ozone formation is dependent on sunlight, the concentration levels vary substantially over the course of the day, and exposure values are therefore affected considerably by what estimation technique is used [90]. We have applied an 8 h maximum as recommended by WHO [8]. Moreover, the same assigned exposure could result in different outcomes depending on how much time a particular individual spends outdoors, under that exposure, thus affecting the actual dose of ozone. The country-based exposures, in the future projections in this study, are based on climate and emission models that utilize a $50 \times 50 \text{ km}$ horizontal resolution. From a continental and country-wide perspective, this resolution can be regarded as high, but on an individual level, it must be seen as rather crude. This introduces a risk of exposure misclassification. For the same reason, ozone exposure is likely overestimated in those areas close to high emission areas, and likewise underestimated in suburban locations, where local emissions could affect the near-surface ozone levels [13].

Thirdly, when including the attributable effects for future air pollution emission levels in EU30, the estimated number of preterm births in 2050 was reduced by 30% when compared to 2010. This is equal to almost 8000 less cases of preterm birth in 2050, meaning that almost one third of all preterm births attributable to ozone exposure could be avoided if ozone precursor emissions follow the predicted emission scenarios. The leading cause behind these drops is the improved technique for reducing vehicular exhaust emissions—e.g. the development of three-way catalytic converters—and legislation regarding ozone precursor emissions [91]. Nevertheless, several uncertainties are related to ozone modelling itself. In our analysis we have applied emission data from the periods of 2005–2010 and 2050 and climate data from the periods of 1991–2000 and

2046–2055. If the future period is centred around 2050, the current climate data is from earlier period than 2010. This might have made bias due to climate change in the last decade [92]. It is also important to note that this estimation does not consider changes in the preterm birth or birth rate as there are no projections on future rates for 2050. During the period of 2010 and 2015 there has been increase of preterm births in some countries and decrease in others [93], but this information does not allow us to make long-term projections. If the preterm birth incidence or birth rate increases/decreases in EU30, the change of attributable cases is misestimated.

Fourthly, when we modelled and projected for changes in the climate, ozone concentration levels for 2050 were found to be relatively similar to those of year 2010, thus leaving the attributable effects relatively unchanged (a 1.6% increased AP in the climate model was seen). The minor impact of ozone exposure in the climate model stems from the fact that the European ozone concentrations did not increase substantially in the current future projections. The highest increase in ozone concentration, due to climate change, was found in Luxembourg, where SOMO25 rose by 12.9% in 2050. If we would calculate the factual daily exposure for this increase, it is equal to a $2.4 \mu\text{g m}^{-3}$ daily increase, which partly can explain the very small effects seen in the climate model.

Apart from the limitations already discussed, this study unavoidably also inherits restrictions from the original studies from which the current exposure-response coefficient has been calculated. All studies in the meta-analysis [63, 64, 69, 71, 74] share the risk of exposure misclassification, as individual exposures are assigned based on ozone concentrations registered by local measurement stations, typically located at rooftop level throughout the city. As previous research, and earlier discussion has indicated [94], this does not, necessarily, reflect the true individual exposure as other meaningful exposure factors, such as time spent outdoors and commute distance, are not taken into account.

When interpreting these findings, one must also consider an excessive degree of uncertainty. Ultimately, it is the exposure-response function which yields the final results of this study, but within the exposure-response function there are factors with varying underlying degrees of uncertainty. The effect size is, for instance, expressed as a relative risk with a 95% confidence interval ranging from 1.009 to 1.046, which puts the average estimate of 27 871 preterm births to be between 9611 and 45 854 cases. Another crucial factor for the function, and subsequently the results, is near-surface ozone concentration. First, we have used 1991–2000 climate data together with 2005–2010 emission data, which may have biased the estimated concentrations. Moreover,

air pollutant emissions are affected by uncertainties in emission factors, which are largely related to real-life end-of-pipe emission control and actual activity data. As shown by Lin *et al* [95] the errors for air pollution emissions could range from 10% to 170% depending on the pollutant and region; however, air pollution emissions are relatively smaller for Western Europe. Many uncertainties are also related to ozone modelling, itself, and atmospheric chemical and physical processes such as dry deposition, wet scavenging, ozone precursors transport, and ozone formation [96, 97]. According to Derwent *et al* [98], the uncertainties from ozone sources and sinks are larger than in model formulation.

Second, there are a multitude of factors impacting the concentration of ozone in the future. Factors impacting the possible future exposure of individuals to ozone in Europe include the pathway for ozone precursor emissions, climate change and the greenhouse gas emission scenario, and sensitivities and parameterizations of the specific climate models and CTMs used to describe the future evolution. There is inherent uncertainty in what path the future ozone exposure evolution will take. The air pollution emission scenarios projected in this project depend on policy makers decisions and the technical solutions available; there are other possible future evolutions.

Watson *et al* [14] compared the impact of climate change, under Current Legislation Emissions (CLE) scenario, and maximally technically feasible reductions emissions modelled by four CTMs. Langner *et al* [44] compared emission change to the impact of climate change on ozone concentrations in two global climate model simulations. Colette *et al* [99] compared a number of different CTM model sensitivities to climate change scenarios. All these studies show that, for a great fraction of the European population, the main contributing factor in ozone concentrations is the pathway of anthropogenic ozone precursor emissions. Thus, future health impacts will be highly dependent on measures taken, and these measures need to be introduced across the northern hemisphere, not only in Europe. Climate change is projected to lead to higher ozone exposure, a so-called climate penalty. Measures to reduce climate change would also act to reduce ozone exposure in most parts of Europe, but the most effective measure is to reduce anthropogenic precursor emissions.

This assessment has focused on the effects of near-surface ozone exposure on preterm birth in the European region exclusively. It should be recognized that the world-wide attributable number of cases of preterm births due to maternal ozone exposure would be substantially higher. This is not only due to the inclusion of larger populations, but in low-income settings, a combination of megacities, poor ozone precursor emission regulations, and a hot climate

with a lot of sunlight will produce ozone levels which are considerably higher than those found in Europe. An example of this can be found in Latin America, as Sao Paulo, Brazil can experience several days during summer where ambient ozone concentrations may exceed $400 \mu\text{g m}^{-3}$ [100], whereas European levels during the summer typically exceed $200 \mu\text{g m}^{-3}$ and are less than $300 \mu\text{g m}^{-3}$, even in extreme summers, like the one experienced in 2003 [101]. This highlights the essentiality of strict global emission legislation, whose need is most pressing in low-income settings, where current regulations in many cases are insufficient.

5. Conclusion

Air pollution is considered the largest environmental threat to human health. Near-surface ozone has been shown to have numerous adverse health effects including facets of reproductive health. According to our meta-analysis, for each $10 \mu\text{g m}^{-3}$ increase in maternal first trimester ozone exposure concentration, the risk for preterm delivery will increase by 2.7%. Subsequently, for 30 European countries, the population AP of preterm births due to first trimester maternal ozone exposure was estimated currently to 7.2%. This proportion equals to a number of 27 871 babies born prematurely in 2010.

Highlighting the importance of emission control and stern legislation regarding ozone precursor emissions, this assessment likewise found that in 2050, with consideration of population change, an estimated number of around 7500 preterm births will be avoided as a result of reduced ozone precursor emissions. Not taking emission and population change into account, due to climate change the ozone-related preterm birth burden might slightly increase by 2050 in Central and Southern Europe, and decrease in Eastern and Northern Europe.

Data availability statement

The data that support the findings of this study are available upon reasonable request from the authors.

Acknowledgments

This work was carried out within the ACCEPTED project, which is supported by the ERA-ENVHEALTH network (Grant Agreement No. 219337), with funding from ANSES, ADEME, BelSPO, UBA and the Swedish EPA. HO work was supported by the Estonian Ministry of Education and Research grant IUT34-17. CA was also supported by the 20172-18 Belmont Forum and BiodivERsA joint call for research proposals under the Biodiv-Scen ERA-Net COFUND programme and with the funding organisations AKA (326328), ANR (ANR-18-EB14-0007), BMBF (KFZ: 01LC1810A), FORMAS

(2018-02434, 2018-02436, 2018-02437, 2018-02438) and MICINN (APCIN: PCI2018-093149). Further support was also given by Nordic WelfAir (Nordforsk Project No. 75007).

Author contributions

JE conceived the study; JE, BF, and HO designed the research; JE and DO performed the meta analyses; HO made maps; JE and HO made the health impact calculations; CA modelled the near-surface ozone and all authors wrote the paper.

Conflict of interest

The authors declare no conflict of interest.

ORCID iD

Hans Orru  <https://orcid.org/0000-0002-7965-9451>

References

- [1] Liu A, Qian N, Yu H, Chen R and Kan H 2017 Estimation of disease burdens on preterm births and low birth weights attributable to maternal fine particulate matter exposure in Shanghai, China *Sci. Total Environ.* **609** 815–21
- [2] Blencowe H, Cousens S, Chou D, Oestergaard M, Say L, Moller A B, Kinney M and Lawn J 2013 Born too soon: the global epidemiology of 15 million preterm births *Reprod. Health* **10** S2
- [3] Institute of Medicine 2007 *Preterm Birth: Causes, Consequences, and Prevention* (Washington, DC: National Academies Press)
- [4] Kajantie E, Osmond C, Barker D J and Eriksson J G 2010 Preterm birth—a risk factor for type 2 diabetes? The Helsinki birth cohort study *Diabetes Care* **33** 2623–5
- [5] Veras M M, De Oliveira Alves N, Fajersztajn L and Saldiva P 2017 Before the first breath: prenatal exposures to air pollution and lung development *Cell Tissue Res.* **367** 445–55
- [6] Moster D, Lie R T and Markestad T 2008 Long-term medical and social consequences of preterm birth *N. Engl. J. Med.* **359** 262–73
- [7] WHO 2016 *Ambient Air Pollution: A Global Assessment of Exposure and Burden of Disease* (Geneva: World Health Organization)
- [8] WHO 2013 *Review of Evidence on Health Aspects of Air Pollution—REVIHAAP Project: Technical Report* (Copenhagen: WHO Regional Office for Europe)
- [9] Toro M V, Cremades L V and Calbó J 2006 Relationship between VOC and NOx emissions and chemical production of tropospheric ozone in the Aburrá Valley (Colombia) *Chemosphere* **65** 881–8
- [10] Jacob D J and Winner D A 2009 Effect of climate change on air quality *Atmos. Environ.* **43** 51–63
- [11] McMichael A J, Woodruff R E and Hales S 2006 Climate change and human health: present and future risks *Lancet* **367** 859–69
- [12] Doherty R M, Heal M R and O'Connor F M 2017 Climate change impacts on human health over Europe through its effect on air quality *Environ. Health: Glob. Access Sci. Source* **16** 118
- [13] Markakis K, Valari M, Engardt M, Lacrosonniere G, Vautard R and Andersson C 2016 Mid-21st century air quality at the urban scale under the influence of changed climate and emissions—case studies for Paris and Stockholm *Atmos. Chem. Phys.* **16** 1877–94
- [14] Watson L et al 2016 Impact of emissions and +2 °C climate change upon future ozone and nitrogen dioxide over Europe *Atmos. Environ.* **142** 271–85
- [15] Krzyzanowski M and Cohen A 2008 Update of WHO air quality guidelines *Air Qual. Atmos. Health* **1** 7–13
- [16] Amann M et al 2008 *Health Risks of Ozone from Long-range Transboundary Air Pollution* (Copenhagen: WHO Regional Office Europe)
- [17] Bell M L, Peng R D and Dominici F 2006 The exposure-response curve for ozone and risk of mortality and the adequacy of current ozone regulations *Environ. Health Perspect.* **114** 532–6
- [18] Chan C C and Wu T H 2005 Effects of ambient ozone exposure on mail carriers' peak expiratory flow rates *Environ. Health Perspect.* **113** 735–8
- [19] Hoek G, Schwartz J D, Groot B and Eilers P 1997 Effects of ambient particulate matter and ozone on daily mortality in Rotterdam, The Netherlands *Arch. Environ. Health* **52** 455–63
- [20] Kim S Y, Lee J T, Hong Y C, Ahn K J and Kim H 2004 Determining the threshold effect of ozone on daily mortality: an analysis of ozone and mortality in Seoul, Korea, 1995–1999 *Environ. Res.* **94** 113–9
- [21] Orru H, Åström C, Andersson C, Tamm T, Ebi K L and Forsberg B 2019 Ozone and heat-related mortality in Europe in 2050 significantly affected by changes in climate, population and greenhouse gas emission *Environ. Res. Lett.* **14** 074013
- [22] Orru H, Ebi K L and Forsberg B 2017 The interplay of climate change and air pollution on health *Curr. Environ. Health Rep.* **4** 504–13
- [23] Chen R, Cai J, Meng X, Kim H, Honda Y, Guo Y L, Samoli E, Yang X and Kan H 2014 Ozone and daily mortality rate in 21 cities of East Asia: how does season modify the association? *Am. J. Epidemiol.* **180** 729–36
- [24] Li T, Yan M, Ma W, Ban J, Liu T, Lin H and Liu Z 2015 Short-term effects of multiple ozone metrics on daily mortality in a megacity of China *Environ. Sci. Pollut. Res. Int.* **22** 8738–46
- [25] Orru H, Andersson C, Ebi K L, Langner J, Åström C and Forsberg B 2013 Impact of climate change on ozone-related mortality and morbidity in Europe *Eur. Respir. J.* **41** 285–94
- [26] Liu Y, Xu J, Chen D, Sun P and Ma X 2019 The association between air pollution and preterm birth and low birth weight in Guangdong, China *BMC Public Health* **19** 3
- [27] Ritz B, Yu F, Fruin S, Chapa G, Shaw G M and Harris J A 2002 Ambient air pollution and risk of birth defects in Southern California *Am. J. Epidemiol.* **155** 17–25
- [28] Salam M T, Millstein J, Li Y F, Lurmann F W, Margolis H G and Gilliland F D 2005 Birth outcomes and prenatal exposure to ozone, carbon monoxide, and particulate matter: results from the children's health study *Environ. Health Perspect.* **113** 1638–44
- [29] Kelly F J 2003 Oxidative stress: its role in air pollution and adverse health effects *Occup. Environ. Med.* **60** 612–6
- [30] Kannan S, Misra D P, Dvonch J T and Krishnakumar A 2006 Exposures to airborne particulate matter and adverse perinatal outcomes: a biologically plausible mechanistic framework for exploring potential effect modification by nutrition *Environ. Health Perspect.* **114** 1636–42
- [31] Shah P S and Balkhair T 2011 Air pollution and birth outcomes: a systematic review *Environ. Int.* **37** 498–516
- [32] Vadillo-Ortega F, Osornio-Vargas A, Buxton M A, Sánchez B N, Rojas-Bracho L, Viveros-Alcaráz M, Castillo-Castrejon M, Beltrán-Montoya J, Brown D G and O'Neill M S 2014 Air pollution, inflammation and preterm birth: a potential mechanistic link *Med. Hypotheses* **82** 219–24
- [33] Catov J M, Bodnar L M, Ness R B, Barron S J and Roberts J M 2007 Inflammation and dyslipidemia related

- to risk of spontaneous preterm birth *Am. J. Epidemiol.* **166** 1312–9
- [34] Lee P C, Talbott E O, Roberts J M, Catov J M, Sharma R K and Ritz B 2011 Particulate air pollution exposure and C-reactive protein during early pregnancy *Epidemiology* **22** 524–31
- [35] Lock K 2000 Health impact assessment *BMJ* **320** 1395–8
- [36] Abe K C and Miraglia S G 2016 Health impact assessment of air pollution in São Paulo, Brazil *Int. J. Environ. Res. Public Health* **13** 694
- [37] Chanel O, Henschel S, Goodman P G, Analitis A, Atkinson R W, Le Tertre A, Zeka A and Medina S 2014 Economic valuation of the mortality benefits of a regulation on SO₂ in 20 European cities *Eur. J. Public Health* **24** 631–7
- [38] Pascal M et al 2013 Assessing the public health impacts of urban air pollution in 25 European cities: results of the Aphekom project *Sci. Total Environ.* **449** 390–400
- [39] Perez L et al 2013 Chronic burden of near-roadway traffic pollution in 10 European cities (APHEKOM network) *Eur. Respir. J.* **42** 594–605
- [40] Malley C S, Kuylenstierna J C, Vallack H W, Henze D K, Blencowe H and Ashmore M R 2017 Preterm birth associated with maternal fine particulate matter exposure: a global, regional and national assessment *Environ. Int.* **101** 173–82
- [41] Andersson C, Bergström R, Bennet C, Robertson L, Thomas M, Korhonen H, Lehtinen K E J and Kokkola H 2015 MATCH-SALSA—multi-scale atmospheric transport and chemistry model coupled to the SALSA aerosol microphysics model—part 1: model description and evaluation *Geosci. Model Dev.* **8** 171–89
- [42] Andersson C, Langner J and Bergström R 2007 Interannual variation and trends in air pollution over Europe due to climate variability during 1958–2001 simulated with a regional CTM coupled to the ERA40 reanalysis *Tellus B* **59** 77–98
- [43] Robertson L, Langner J and Engardt M 1999 An eulerian limited-area atmospheric transport model *J. Appl. Meteorol.* **38** 190–210
- [44] Langner J, Bergström R and Pleijel K 1998 European scale modeling of sulfur, oxidised nitrogen and photochemical oxidants. Model development and evaluation for the 1994 growing season *RMK No 82* (Swedish Meteorological and Hydrological Institute)
- [45] Simpson D, Guenther A, Hewitt C N and Steinbrecher R 1995 Biogenic emissions in Europe: 1. Estimates and uncertainties *J. Geophys. Res.: Atmos.* **100** 22875–90
- [46] Carter W P L 1996 Condensed atmospheric photooxidation mechanisms for isoprene *Atmos. Environ.* **30** 4275–90
- [47] Simpson D et al 2012 The EMEP MSC-W chemical transport model—technical description *Atmos. Chem. Phys.* **12** 7825–65
- [48] Atkinson R, Baulch D L, Cox R A, Crowley J N, Hampson R F, Hynes R G, Jenkin M E, Rossi M J, Troe J and Subcommittee I 2006 Evaluated kinetic and photochemical data for atmospheric chemistry: volume II—gas phase reactions of organic species *Atmos. Chem. Phys.* **6** 3625–4055
- [49] Andersson C and Engardt M 2010 European ozone in a future climate: importance of changes in dry deposition and isoprene emissions *J. Geophys. Res.: Atmos.* **115** D02303
- [50] Marécal V et al 2015 A regional air quality forecasting system over Europe: the MACC-II daily ensemble production *Geosci. Model Dev.* **8** 2777–813
- [51] Andersson C, Alpfjör H, Robertson L, Karlsson P E and Engardt M 2017 Reanalysis of and attribution to near-surface ozone concentrations in Sweden during 1990–2013 *Atmos. Chem. Phys.* **17** 13869–90
- [52] Otero N et al 2018 A multi-model comparison of meteorological drivers of surface ozone over Europe *Atmos. Chem. Phys.* **18** 12269–88
- [53] Lacressonnière G et al 2016 Impacts of regional climate change on air quality projections and associated uncertainties *Clim. Change* **136** 309–24
- [54] Watson L et al 2015 The impact of meteorological forcings on gas phase air pollutants over Europe *Atmos. Environ.* **119** 240–57
- [55] Strandberg G, Bärning L, Hansson U, Jansson C, Jones C, Kjellström E, Kupiainen M, Nikulin G, Samuelsson P and Ullerstig A 2015 CORDEX scenarios for Europe from the Rossby centre regional climate model *RCA4 Report Meteorology and Climatology No. 116* 2014 (Norrköping: SMHI)
- [56] Thomson A M et al 2011 RCP4.5: a pathway for stabilization of radiative forcing by 2100 *Clim. Change* **109** 77
- [57] Hauglustaine D A, Hourdin F, Jourdain L, Filiberti M-A, Walters S, Lamarque J-F and Holland E A 2004 Interactive chemistry in the Laboratoire de Météorologie Dynamique general circulation model: description and background tropospheric chemistry evaluation *J. Geophys. Res.: Atmos.* **109** D04314
- [58] UN 2020 Preterm birth rate (per 100 live births) (United Nations Statistics Division) (available at: [http://data.un.org/Data.aspx?d=WHO&f=MEASURE_CODE%3aWHS_PBR:](http://data.un.org/Data.aspx?d=WHO&f=MEASURE_CODE%3aWHS_PBR;))
- [59] WHO 2020 Global health observatory visualizations, indicator metadata registry (World Health Organization) (available at: <https://apps.who.int/gho/data/node.wrapper.imr?x-id=1:>)
- [60] Eurostat 2019 EUROPOP2019—population projections at national level (2019–2100) ed Eurostat
- [61] Arroyo V, Díaz J, Carmona R, Ortiz C and Linares C 2016 Impact of air pollution and temperature on adverse birth outcomes: Madrid, 2001–2009 *Environ. Pollut.* **218** 1154–61
- [62] Capobussi M et al 2016 Air pollution impact on pregnancy outcomes in Como, Italy *J. Occup. Environ. Med.* **58** 47–52
- [63] Ha S, Hu H, Roussos-Ross D, Haidong K, Roth J and Xu X 2014 The effects of air pollution on adverse birth outcomes *Environ. Res.* **134** 198–204
- [64] Hansen C, Neller A, Williams G and Simpson R 2006 Maternal exposure to low levels of ambient air pollution and preterm birth in Brisbane, Australia *BJOG: Int. J. Obstetrics Gynaecol.* **113** 935–41
- [65] Jalaludin B, Mannes T, Morgan G, Lincoln D, Sheppard V and Corbett S 2007 Impact of ambient air pollution on gestational age is modified by season in Sydney, Australia *Environ. Health: Glob. Access Science Source* **6** 16
- [66] Laurent O, Hu J, Li L, Kleeman M J, Bartell S M, Cockburn M, Escobedo L and Wu J 2016 A statewide nested case-control study of preterm birth and air pollution by source and composition: California, 2001–2008 *Environ. Health Perspect.* **124** 1479–86
- [67] Lavigne E et al 2016 Ambient air pollution and adverse birth outcomes: differences by maternal comorbidities *Environ. Res.* **148** 457–66
- [68] Lee P C, Roberts J M, Catov J M, Talbott E O and Ritz B 2013 First trimester exposure to ambient air pollution, pregnancy complications and adverse birth outcomes in Allegheny County, PA *Maternal Child Health J.* **17** 545–55
- [69] Lin Y T, Jung C R, Lee Y L and Hwang B F 2015 Associations between ozone and preterm birth in women who develop gestational diabetes *Am. J. Epidemiol.* **181** 280–7
- [70] Liu S, Krewski D, Shi Y, Chen Y and Burnett R T 2003 Association between gaseous ambient air pollutants and adverse pregnancy outcomes in Vancouver, Canada *Environ. Health Perspect.* **111** 1773–8
- [71] Olsson D, Ekström M and Forsberg B 2012 Temporal variation in air pollution concentrations and preterm

- birth—a population based epidemiological study *Int. J. Environ. Res. Public Health* **9** 272–85
- [72] Qian Z *et al* 2016 Ambient air pollution and preterm birth: a prospective birth cohort study in Wuhan, China *Int. J. Hyg. Environ. Health* **219** 195–203
- [73] Ritz B, Yu F, Chapa G and Fruin S 2000 Effect of air pollution on preterm birth among children born in Southern California between 1989 and 1993 *Epidemiology* **11** 502–11
- [74] Olsson D, Mogren I and Forsberg B 2013 Air pollution exposure in early pregnancy and adverse pregnancy outcomes: a register-based cohort study *BMJ Open* **3** e001955
- [75] Harris R, Bradburn M, Deeks J, Harbord R, Altman D, Steichen T and Sterne J 2006 METAN: stata module for fixed and random effects meta-analysis *Stat. Softw. Compon.* S456798
- [76] Borenstein M, Hedges L V, Higgins J P and Rothstein H R 2010 A basic introduction to fixed-effect and random-effects models for meta-analysis *Res. Synth. Methods* **1** 97–111
- [77] DerSimonian R and Laird N 1986 Meta-analysis in clinical trials *Control Clin. Trials* **7** 177–88
- [78] Sun X, Luo X, Zhao C, Chung Ng R W, Lim C E, Zhang B and Liu T 2015 The association between fine particulate matter exposure during pregnancy and preterm birth: a meta-analysis *BMC Pregnancy Childbirth* **15** 300
- [79] Bosetti C, Nieuwenhuijsen M J, Gallus S, Cipriani S, La Vecchia C and Parazzini F 2010 Ambient particulate matter and preterm birth or birth weight: a review of the literature *Arch. Toxicol.* **84** 447–60
- [80] Stieb D M, Chen L, Eshoul M and Judek S 2012 Ambient air pollution, birth weight and preterm birth: a systematic review and meta-analysis *Environ. Res.* **117** 100–11
- [81] Menon R 2008 Spontaneous preterm birth, a clinical dilemma: etiologic, pathophysiologic and genetic heterogeneities and racial disparity *Acta Obstet. Gynecol. Scand.* **87** 590–600
- [82] Chen G, Guo Y, Abramson M J, Williams G and Li S 2018 Exposure to low concentrations of air pollutants and adverse birth outcomes in Brisbane, Australia, 2003–2013 *Sci. Total Environ.* **622–623** 721–6
- [83] Dastoorpoor M, Idani E, Goudarzi G and Khanjani N 2018 Acute effects of air pollution on spontaneous abortion, premature delivery, and stillbirth in Ahvaz, Iran: a time-series study *Environ. Sci. Pollut. Res. Int.* **25** 5447–58
- [84] Fleischer N L, Merialdi M, Van Donkelaar A, Vadillo-Ortega F, Martin R V, Betran A P and Souza J P 2014 Outdoor air pollution, preterm birth, and low birth weight: analysis of the world health organization global survey on maternal and perinatal health *Environ. Health Perspect.* **122** 425–30
- [85] Li X *et al* 2017 Association between ambient fine particulate matter and preterm birth or term low birth weight: an updated systematic review and meta-analysis *Environ. Pollut.* **227** 596–605
- [86] UNECE 2003 *Modelling and Assessment of the Health Impact of Particulate Matter and Ozone: Summary Report* (Geneva: UN)
- [87] Anderson H R, Atkinson R W, Peacock J, Marston L, Konstantinou K and Organization W H 2004 *Meta-analysis of Time-series Studies and Panel Studies of Particulate Matter (PM) and Ozone (O₃): Report of a WHO Task Group* (Copenhagen: WHO Regional Office for Europe)
- [88] De Marco A 2009 Assessment of present and future risk to Italian forests and human health: modelling and mapping *Environ. Pollut.* **157** 1407–12
- [89] Mitis F, Iavarone I and Martuzzi M 2007 Health impact of ozone in 13 Italian cities *Epidemiol. Prev.* **31** 323–32
- [90] Bell M and Ellis H 2003 Comparison of the 1 hr and 8 hr national ambient air quality standards for ozone using models-3 *J. Air Waste Manage. Assoc.* **53** 1531–40
- [91] EEA 2015 Emissions of ozone precursors (European Environment Agency) (available at: www.eea.europa.eu/data-and-maps/indicators/emissions-of-ozone-precursors-version-2/assessment-4.)
- [92] WMO 2013 The global climate 2001–2010: a decade of climate extremes *Summary Report* (Geneva: World Meteorological Organization)
- [93] Euro-Peristat Project 2018 European perinatal health report Core indicators of the health and care of pregnant women and babies in Europe in 2015
- [94] Liu L J, Koutrakis P, Suh H H, Mulik J D and Burton R M 1993 Use of personal measurements for ozone exposure assessment: a pilot study *Environ. Health Perspect.* **101** 318–24
- [95] Lin J *et al* 2019 Carbon and health implications of trade restrictions *Nat. Commun.* **10** 4947
- [96] Dunker A M, Wilson G, Bates J T and Yarwood G 2020 Chemical sensitivity analysis and uncertainty analysis of ozone production in the comprehensive air quality model with extensions applied to Eastern Texas *Environ. Sci. Technol.* **54** 5391–9
- [97] Boynard A, Beekmann M, Foret G, Ung A, Szopa S, Schmechtig C and Coman A 2011 An ensemble assessment of regional ozone model uncertainty with an explicit error representation *Atmos. Environ.* **45** 784–93
- [98] Derwent R G, Parrish D D, Galbally I E, Stevenson D S, Doherty R M, Naik V and Young P J 2018 Uncertainties in models of tropospheric ozone based on Monte Carlo analysis: tropospheric ozone burdens, atmospheric lifetimes and surface distributions *Atmos. Environ.* **180** 93–102
- [99] Colette A *et al* 2015 Is the ozone climate penalty robust in Europe? *Environ. Res. Lett.* **10** 084015
- [100] WHO 2006 *Air Quality Guidelines: Global Update 2005* (Copenhagen: WHO Regional Office for Europe)
- [101] Solberg S, Ø H, Søvde A, Isaksen I S A, Coddeville P, De Backer H, Forster C, Orsolini Y and Uhse K 2008 European surface ozone in the extreme summer 2003 *J. Geophys. Res.: Atmos.* **113** D07307