

The coupled hydrology–human activity information (CHHAI) dataset: a global benchmark dataset for model comparison and evaluation

Azara Boschee, Aneseh Alborzi, Augustina Clara Alexander, Berit Arheimer, Salsebil Bel Hadj Ali, Günter Blöschl, Andrea Castelletti, Xi Chen, Aniya Dogra, Erhu Du, Jesse Duku, Aldo Fiori, Margaret Garcia, Salvatore Grimaldi, Naota Hanasaki, Yeonjoo Kim, Heidi Kreibich, Chenyuan Li, Mir Matin, Pedro Medeiros, Ali Mehran, Antônio Alves Meira Neto, Fuko Nakai, Shinichiro Nakamura, Phu Nguyen, Joel Nobert, Afua Owusu, Saritha Padiyedath Gopalan, Bruno Pereira, Parsa Pouladi, Mehrsa Pouladi, Mahendran Roobavannan, Mojtaba Sadegh, Matteo Sangiorgio, Lukas Schoppa, Ashish Shrestha, Murugesu Sivapalan, Deborah Sousa, Sai Veena Sunkara, Fuqiang Tian, Fatma Trabelsi, Naga Velpuri, Elena Volpi, Jiale Wang, Shuo Wang, Jiefeng Wu, Xing Yuan & Amir AghaKouchak

To cite this article: Azara Boschee, Aneseh Alborzi, Augustina Clara Alexander, Berit Arheimer, Salsebil Bel Hadj Ali, Günter Blöschl, Andrea Castelletti, Xi Chen, Aniya Dogra, Erhu Du, Jesse Duku, Aldo Fiori, Margaret Garcia, Salvatore Grimaldi, Naota Hanasaki, Yeonjoo Kim, Heidi Kreibich, Chenyuan Li, Mir Matin, Pedro Medeiros, Ali Mehran, Antônio Alves Meira Neto, Fuko Nakai, Shinichiro Nakamura, Phu Nguyen, Joel Nobert, Afua Owusu, Saritha Padiyedath Gopalan, Bruno Pereira, Parsa Pouladi, Mehrsa Pouladi, Mahendran Roobavannan, Mojtaba Sadegh, Matteo Sangiorgio, Lukas Schoppa, Ashish Shrestha, Murugesu Sivapalan, Deborah Sousa, Sai Veena Sunkara, Fuqiang Tian, Fatma Trabelsi, Naga Velpuri, Elena Volpi, Jiale Wang, Shuo Wang, Jiefeng Wu, Xing Yuan & Amir AghaKouchak (04 Jun 2026): The coupled hydrology–human activity information (CHHAI) dataset: a global benchmark dataset for model comparison and evaluation, Hydrological Sciences Journal, DOI: [10.1080/02626667.2026.2663050](https://doi.org/10.1080/02626667.2026.2663050)

To link to this article: <https://doi.org/10.1080/02626667.2026.2663050>



© 2026 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 04 Jun 2026.



Submit your article to this journal [↗](#)



Article views: 778



[View related articles](#) 



[View Crossmark data](#) 

The coupled hydrology–human activity information (CHHA) dataset: a global benchmark dataset for model comparison and evaluation

Azara Boschee^a, Aneseh Alborzi^a, Augustina Clara Alexander^b, Berit Arheimer^c, Salsebil Bel Hadj Ali^d, Günter Blöschl^{e,f}, Andrea Castelletti^{g,h}, Xi Chenⁱ, Aniya Dogra^j, Erhu Du^k, Jesse Duku^l, Aldo Fiori^l, Margaret Garcia^m, Salvatore Grimaldiⁿ, Naota Hanasaki^{o,p}, Yeonjoo Kim^q, Heidi Kreibich^r, Chenyuan Li^s, Mir Matin^t, Pedro Medeiros^u, Ali Mehran^j, Antônio Alves Meira Neto^v, Fuko Nakai^w, Shinichiro Nakamura^x, Phu Nguyen^a, Joel Nobert^y, Afua Owusu^z, Saritha Padiyedath Gopalan^o, Bruno Pereira^{aa}, Parsa Pouladi^{bb,cc}, Mehrsa Pouladi^{dd}, Mahendran Roobavannan^{ee}, Mojtaba Sadegh^{ff}, Matteo Sangiorgio^g, Lukas Schoppa^{h,gg}, Ashish Shrestha^{hh}, Murugesu Sivapalanⁱⁱ, Deborah Sousa^{aa}, Sai Veena Sunkara^{jj}, Fuqiang Tian^{kk}, Fatma Trabelsi^d, Naga Velpuri^l, Elena Volpi^l, Jiale Wang^{kk}, Shuo Wang^{mm}, Jiefeng Wuⁿⁿ, Xing Yuan^s and Amir AghaKouchak^l

^aDepartment of Civil and Environmental Engineering, University of California, Irvine, California, USA; ^bDepartment of Water Resources Engineering – CoET, University of Dar es Salaam, Dar es Salaam, Tanzania; ^cDepartment of Hydrological Research, Swedish Meteorological and Hydrological Institute (SMHI), Norrköping, Sweden; ^dResearch Unit Sustainable Management of Water and Soil Resources, Higher School of Engineers of Medjeb El Bab (ESIM), University of Jendouba, Jendouba, Tunisia; ^eInstitute of Hydraulic Engineering and Water Resources Management, Vienna University of Technology, Vienna, Austria; ^fDepartment of Civil, Chemical Environmental and Materials Engineering, University of Bologna, Bologna, Italy; ^gDepartment of Electronics, Information and Bioengineering, Politecnico di Milano, Milano, Italy; ^hCMCC Foundation – Euro-Mediterranean Centre on Climate Change, Milano, Italy; ⁱDepartment of Geography and Geographic Information Science, University of Cincinnati, Cincinnati, Ohio, USA; ^jDepartment of Civil and Environmental Engineering, San Jose State University, San Jose, California, USA; ^kYangtze Institute for Conservation and Development, Hohai University, Nanjing, China; ^lDepartment of Civil, Computer Science, and Aeronautical Technologies Engineering, Roma Tre University, Rome, Italy; ^mSchool of Sustainable Engineering and the Built Environment, Arizona State University, Tempe, Arizona, USA; ⁿDIBAF Department, Tuscia University, Viterbo, Italy; ^oDepartment of Civil Engineering, The University of Tokyo, Tokyo, Japan; ^pNational Institute for Environmental Studies (NIES), Tsukuba, Japan; ^qDepartment of Civil and Environmental Engineering, Yonsei University, Seoul, South Korea; ^rSection Hydrology, GFZ Helmholtz Centre for Geosciences, Potsdam, Germany; ^sState Key Laboratory of Earth System Numerical Modeling and Application, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China; ^tUnited Nations University Institute for Water, Environment, and Health (UNU-INWEH), Richmond Hill, ON, Canada; ^uDepartment of Civil Engineering, Federal Institute of Education, Science and Technology of Ceará, Fortaleza, Brazil; ^vDepartment of Civil and Environmental Engineering, Colorado State University, Fort Collins, Colorado, USA; ^wCenter for Spatial Information Science, The University of Tokyo, Tokyo, Japan; ^xDepartment of Civil and Environmental Engineering, Nagoya University, Nagoya, Japan; ^yDepartment of Water Resources Engineering, University of Dar es Salaam, Dar es Salaam, Tanzania; ^zInternational Water Management Institute, Accra, Ghana; ^{aa}Environmental Technology and Water Resources Graduate Program, University of Brasília, Brasília, Brazil; ^{bb}Lyles School of Civil Engineering, Purdue University, West Lafayette, Indiana, USA; ^{cc}Ecological Sciences and Engineering Interdisciplinary Graduate Program, Purdue University, West Lafayette, Indiana, USA; ^{dd}Department of Civil Engineering, Shahid Beheshti University, Tehran, Iran; ^{ee}School of Civil and Environmental Engineering, Faculty of Engineering and Information Technology, University of Technology, Sydney, Australia; ^{ff}Department of Civil Engineering, Boise State University, Boise, Idaho, USA; ^{gg}Munich Reinsurance Company, Munich, Germany; ^{hh}Urban Systems Lab, New York University, New York, New York, USA; ⁱⁱDepartment of Civil and Environmental Engineering, University of Illinois, Urbana-Champaign, Urbana, Illinois, USA; ^{jj}School of Civil and Environmental Engineering, Cornell University, Ithaca, New York, USA; ^{kk}Department of Hydraulic Engineering, State Key Laboratory of Hydrosience and Engineering, Tsinghua University, Beijing, China; ^{ll}International Water Management Institute, Colombo, Sri Lanka; ^{mm}Department of Land Surveying and Geo-Informatics, The Hong Kong Polytechnic University, Hung Hom, Kowloon, Hong Kong; ⁿⁿCollege of Civil Engineering, Hefei University of Technology, Hefei, China

ABSTRACT

To enhance hydrologic modeling, the hydrology community has developed benchmark datasets (e.g. Model Parameter Estimation Experiment, MOPEX), providing standardized data for model evaluation and parameter estimation. However, these datasets primarily focus on modeling natural hydrologic processes, leaving a critical gap in understanding the role of human influences. Here, we introduce the Coupled Hydrology-Human Activity Information (CHHA) dataset, a benchmark dataset that integrates coupled human–water data from regions across all continents, excluding Antarctica. CHHA incorporates data from 25 regions that cover various human impacts such as reservoir management, flood protection, river management policies, land use changes, and water use. Each basin reflects distinct challenges, providing a diverse and globally representative resource for researchers studying these processes. By offering standardized datasets for modeling and analysis, CHHA aims to enhance our understanding of interactions between people and water and to support the development of improved strategies for managing coupled human–water systems.

ARTICLE HISTORY

Received 18 September 2025
Accepted 20 March 2026

EDITOR

A. Castellarin

ASSOCIATE EDITOR

K. Schröter

KEYWORDS

Coupled human–water systems; sociohydrology; hydrologic data; sociohydrologic modeling

Introduction

The hydrologic modeling community has long recognized the importance of standardized benchmark datasets for calibration, validation, and evaluating model performance. Notable examples include the Model Parameter Estimation Experiment (MOPEX; Duan *et al.* 2006, Schaake *et al.* 2006) and the data provided as part of the Distributed Model Intercomparison Project (DMIP; Reed *et al.* 2004, Smith *et al.* 2013), which have facilitated systematic evaluations of hydrologic models across diverse natural basins. Building on MOPEX, the Catchment Attributes and MEteorology for Large-Sample Studies (CAMELS; Newman *et al.* 2015, Addor *et al.* 2017) dataset provides daily forcing data and streamflow for small to medium-sized watersheds in the United States. This led to efforts to provide CAMELS data for basins in other countries such as Chile (AlvarezGarretón *et al.* 2018), Great Britain (Coxon *et al.* 2020), and Brazil (Chagas *et al.* 2020). More recently, Caravan (Kratzert *et al.* 2023) standardized and aggregated the CAMELS data into a unified database while EStreams (Do Nascimento *et al.* 2024) integrated catchment data across Europe into a single dataset. Although these datasets provide widespread and long-term hydrological data, they have predominantly focused on natural hydrologic systems (Addor *et al.* 2020) or effects of climate change (Turner *et al.* 2025), largely excluding the direct influence of human activities (e.g. water withdrawals, reservoir operations, irrigation practices, and interbasin water transfers). Related efforts have begun to incorporate sociohydrologic information, including the Panta Rhei benchmark dataset, which consists of sociohydrological data on paired flood and drought events (Kreibich *et al.* 2023).

In the past few decades, particularly during the International Association of Hydrological Sciences (IAHS) scientific decade Panta Rhei – Change in hydrology and society (2013–2022; Montanari *et al.* 2013, Kreibich *et al.* 2025, Tian *et al.* 2026), the hydrology community has increasingly recognized the significant influence of human activities on the water cycle, including their role in altering streamflow patterns, groundwater dynamics, and the frequency and intensity of hydrologic extremes such as floods and droughts (Oki and Kanae 2006, Peel and Blöschl 2011, Sivapalan and Blöschl 2015, Troy *et al.* 2015b, Blair and Buytaert 2016, AghaKouchak *et al.* 2021, Kreibich *et al.* 2022). As a result, a new generation of regional and global hydrologic models has emerged, explicitly incorporating human interventions such as water withdrawals, reservoir operations, irrigation practices, interbasin transfers, and urban water use. Models developed to simulate these interactions include the Water Global Assessment and Prognosis (WaterGAP; Müller Schmied *et al.* 2021), the Water Balance Model (WBM; Wisser *et al.* 2008, Grogan *et al.* 2022), H08 (Hanasaki *et al.* 2018), the PCRaster Global Water Balance model (PCRGLOBWB; van Beek and Bierkens 2008, Sutanudjaja *et al.* 2018) and the WorldWide HYPE model (Arheimer *et al.* 2020). These models incorporate human water demand to simulate water storage, whereas the Water Evaluation and Planning System (WEAP; Sieber 2006) and the Community Water Model (CWatM; Burek *et al.* 2020) were developed to assist

with water resource management. Additionally, software packages have been developed to integrate human activities into more traditional hydrological models such as VICRes (Dang *et al.* 2020) which provides tools for representing reservoirs in the Variable Infiltration Capacity model (VIC; Liang *et al.* 1994). Similarly, the widely used Soil and Water Assessment Tool (SWAT; Arnold *et al.* 1998) was created to assist water resource managers in evaluating both water supply and water quality, particularly related to agricultural pollution in watersheds. These human-integrated models aim to better reflect real-world conditions and support more representative assessments of water availability, risks, and management strategies under both current and future scenarios influenced by climate change and socioeconomic development (e.g. Bartosova *et al.* 2019). More recently, machine learning and data science have begun to further enhance hydrologic modeling in human-dominated systems (Maity *et al.* 2024). This makes it even more important to have standardized benchmark datasets tailored for evaluating and comparing the performance of such models in human-dominated systems. In the context of the Panta Rhei decadal initiative, this data challenge is a major feature of the synthesis effort that was recently completed (Kreibich *et al.* 2023, 2025, Tian *et al.* 2026).

Studies that explicitly incorporate bidirectional interactions between water systems and human activities are often referred to as sociohydrology (Sivapalan *et al.* 2012, Troy *et al.* 2015a, Blair and Buytaert 2016, Pande and Sivapalan 2017, Pouladi *et al.* 2025b, Tian *et al.* 2026) or, more broadly, as coupled human–water systems research (Li *et al.* 2017b, Wada *et al.* 2017, Konar *et al.* 2019, Reed *et al.* 2022). Regarding the most appropriate terminology (e.g. reflecting differences in disciplinary focus, modeling approaches, and conceptual frameworks), this study uses both terms interchangeably. Our primary focus is not on the specific classification of these models, but rather on addressing the critical need for benchmark datasets that can support the development, calibration, validation, and intercomparison of hydrologic models that account for human influences on water systems.

A wide range of human activities can alter the hydrologic cycle and influence natural hazard dynamics (Kreibich *et al.* 2025). These interventions not only modify the timing and magnitude of water flows but also affect how we observe, model, and manage hydrologic processes and associated risks. These influences vary between regions and cities due to differences in local needs and government policies (Salvadore *et al.* 2015, Pouladi *et al.* 2025b). Historically, people have settled near water bodies, especially rivers, making them particularly vulnerable to flooding (Kummu *et al.* 2011). In response to flooding events, communities often build levees and relocate inland to mitigate impacts. However, when memory of past floods fades, a false sense of security emerges, prompting renewed settlement closer to rivers, resulting in less frequent but more devastating flood events (Di Baldassarre *et al.* 2013). This phenomenon is termed the levee effect or safe development paradox and is one socio-hydrological phenomenon among many (see, for instance, Di Baldassarre *et al.* 2019 for a list of seven relevant phenomena

and 11 sub-phenomena). In contrast, populations are drawn closer to water sources during droughts (Ceola *et al.* 2023). These patterns reflect the ideas of social memory and risk awareness (Albertini *et al.* 2020).

Land cover changes can influence the water cycle. For example, urban land expansion increases impervious surfaces, leading to increased runoff and flood risk (Rose and Peters 2001, Sui and Van de Ven 2023). Similarly, expansions of agricultural land can potentially increase runoff; however, this effect depends on the characteristics of the fields (Fiener *et al.* 2011). Agricultural land can also lead to higher evapotranspiration due to greater crop coverage (Liu *et al.* 2008, Odongo *et al.* 2019), requiring the integration of crop models in hydrological modeling to improve estimations of leaf area indices and evapotranspiration rates (Zhang *et al.* 2021). Land cover change can also be reversed as seen in cases of the “pendulum swing,” where systems are modified to improve human quality of life (e.g. irrigation systems) before restoration efforts return them to more natural conditions (Kandasamy *et al.* 2014). Conversely, hydrological systems can drive land cover change. For example, flood zone regulations can restrict development to certain areas, indirectly affecting runoff (Pan *et al.* 2018).

As water is a vital resource for consumption, irrigation, and hydropower generation, hydrologic modeling is critical for reservoir management. Reservoir operations substantially influence water availability by regulating streamflow and affecting upstream and downstream water supply, and because they are governed by human decision making, dam construction and reservoir releases can complicate hydrological predictions downstream (Mehran *et al.* 2017, Meira Neto *et al.* 2024). In transboundary river basins, these complications are amplified because reservoir systems are managed by multiple actors, creating feedback loops that require a comprehensive socio-hydrological perspective to inform sustainable water management (Shrestha *et al.* 2022). Given these examples of human interventions in hydrologic systems, appropriate modeling approaches depend on the types of human processes specific to each region. Historically, human variables have been considered boundary conditions (Sivapalan and Blöschl 2015); however, this assumption has been deemed incorrect, as human activities can change over time, meaning that adequate data is necessary to accurately predict coupled human–water systems (Peel and Blöschl 2011).

Despite the well-documented importance of human activities in shaping the hydrologic cycle and the growing body of literature on models and tools that account for these interactions, there remains a critical gap: the absence of a standardized benchmark dataset, similar to MOPEX, for evaluating and comparing the performance of models in human-influenced systems. This need is underscored by the fact that models tend to perform worse in highly regulated areas or when human impacts are not modeled (Tijerina-Kreuzer *et al.* 2021, Mai *et al.* 2022, Meira Neto *et al.* 2024). To address this gap, we introduce the Coupled Hydrology-Human Activity Information dataset (CHHAI). CHHAI integrates coupled human–water data from diverse regions across all continents, excluding Antarctica. The variables and types of data capture a wide range of water–human challenges for each region, leading to a diverse and globally representative

resource that supports context-specific analyses in case study research (Mostert 2018). The dataset can be used to reveal insights into how human impacts are affecting regions (e.g. land use change, reservoir management, water withdrawal for irrigation and use) and how communities are responding to them (e.g. flood protection, the pendulum swing, water demand management).

This effort is inspired by the second and third IAHS scientific decades, spanning 2013–2022 and 2023–2032. The second decade was titled *Panta Rhei – Everything Flows*; it was dedicated to understanding the changes in both hydrology and society as a coupled system (Montanari *et al.* 2013). The third decade, titled *Hydrology Engaging Local People IN one Global world (HELPING)*, has a focus on the use of open science and local data to solve local water problems (Arheimer *et al.* 2024). By focusing on processes and challenges specific to each basin, this effort enables researchers to study localized processes while also identifying global patterns in human–water interactions.

Data and methods

CHHAI contains data for 25 regions around the world, including river and lake basins. To retrieve this data, we contacted researchers studying these regions and invited them to contribute to CHHAI. To be included, data must contain both human-related and surface hydrology data. The types of human-related data that would be accepted are broad, including water use and demand, irrigation and agriculture systems, and reservoir implementation and management. We preferred data with long periods of record; however, if a significant hydrologic or anthropogenic event occurred during the study period, a shorter record of data was accepted. Because each region has unique challenges and scenarios, different data variables and types were accepted; however, most basins have time-series data.

Information about how data for individual basins were collected is described in the papers associated with the data (see Table A1 in the Appendix or the ReadMe files). Once the data were collected, datasets were standardized to have common units (metric system) and a standard format. Time-series data are in the format required for submission to HydroShare, which is maintained by the Consortium of Universities for the Advancement of Hydrologic Science, Inc. (CUAHSI 2025; Horsburgh *et al.* 2016). Geospatial data are shared in either NetCDF or TIFF files, and non-formatted information about a region is provided in tables or in the ReadMe file.

The locations of 25 initial study river/lake basins and non-basin regions included in CHHAI are shown in Fig. 1. The majority of the basins and regions are located in Asia, with at least two regions in other continents. The initial regions exhibit different human impacts that can be studied with the data, such as reservoir management, flood protection, river management policies, land use changes, and water use and withdrawal. It should be noted that the human impacts shown in Fig. 1 do not completely cover all human impacts experienced by a region; rather, they cover only those represented by the data in CHHAI. More characteristics (e.g. data types and variables) of the regional datasets are listed




Figure 1. Locations of the 25 initial river basins, lake basins, and regions included in CHHAI. The basins are colored by the types of major human impacts included in the data for the basin. Cross-hatched colors indicate that multiple types of impacts are reflected in the basin's data. Labels A through E show the locations of the basins that are used in the "Data usage and example analysis" section (Figs 2 through 6, respectively). Map insets highlight boundaries of smaller basins for easier visibility.

in Table A1. The CHHAI dataset is available at <https://doi.org/10.4211/hs.e011b48fabd2465eb5135275d2f3a93e>. Data are organized as separate ZIP folders for each region (see column 6 of Table A1). The ZIP files include a ReadMe file that describes the contents of each folder.

Data usage and example analysis

This section presents example analyses for five basins to illustrate potential research questions that could be investigated using CHHAI. The examples focus on water and




Figure 2. Observed streamflow at the Calpella gauge and average daily maximum allowable water withdrawal (water rights) for the East Fork Russian River in the United States (Label A in Fig. 1). The dashed blue line is the yearly total observed streamflow, and the solid blue line is the 5-year rolling average of the yearly streamflow. For further information on this data, see Garcia *et al.* (2024).

basin management policies and agricultural irrigation practices.

The first example is the East Fork Russian River in California, USA (label “A” in Fig. 1). This river is regulated by Lake Mendocino (formed by the Coyote Valley dam) and the Potter Valley Project, which diverts water from the Eel River to the East Fork Russian River (Garcia *et al.* 2024). In California, the daily maximum amount of allowable water withdrawal (water rights) has increased over time while streamflow in the river has experienced a slight decrease (Fig. 2). From 1979 to 1985, water rights nearly quadrupled (from approximately 0.6 mm/day to 2.6 mm/day). Before 1985, the mean observed streamflow was 1286 mm/year, and after 1985, it decreased to 991 mm/year, a reduction of almost 300 mm/year. This decrease in streamflow can be explained in part by a decline in imported water from the Potter Valley Project; however, the increase in allowed water withdrawals may also have contributed to the reduced streamflow. As there may be other causes of decreased streamflow, future researchers could investigate questions such as the following: What other changes in the watershed contributed to the streamflow decline? How did irrigators respond to changes in streamflow? As the Potter Valley Project is planned to be decommissioned (Alexander 2025), what are the potential impacts resulting from reduced water diverted into the basin? Why have water rights increased over time?

In another example, the Kissimmee River in central Florida, USA (label “B” in Fig. 1) experienced changing river management practices. In 1962, water managers started channelizing the downstream part of the river due to repeated flooding. However, this channelization was found to have negatively affected natural wetlands, and planning began to restore the river to natural conditions (Chen *et al.* 2016). During this period, there were no noticeable changes in streamflow alone in the river (Fig. 3); however, further analysis incorporating land cover and socioeconomic variables may indicate whether channelization did impact river flow. This basin is an example of the “pendulum swing” (Kandasamy *et al.* 2014, Chen *et al.* 2016), where artificial management of a river basin is reversed

to return the river to a more natural state. Potential future research questions include the following: How do different river management strategies (e.g. channelization versus restoration) influence long-term streamflow dynamics and variability? To what extent do restoration efforts reverse the hydrological and ecological impacts of earlier channelization?

Similar to the water withdrawal policies in the Russian River basin, the Tarim River Basin Authority implemented water conservation policies following changes in water use and irrigated areas in the Tarim River basin in China (label “C” in Fig. 1). The Tarim River basin has been widely recognized as a canonical sociohydrological system, where irrigation expansion, ecological degradation, and policy interventions coevolved over multiple decades (Liu *et al.* 2014, Ye *et al.* 2014, 2015). Throughout the late 1990s, the Tarim River basin experienced increasing farmland and irrigation infrastructure. In response, the Tarim River Basin Authority initiated water conservation projects to constrain water use for irrigation and emergency water transfers. These transfers allocated water to downstream lakes in order to preserve vegetation (Liu *et al.* 2014). These changes, along with the resulting upstream and downstream streamflow in the Tarim river, can be seen in Fig. 4. The average trends in upstream and downstream streamflow are -7.94 million cubic meters per year and -30.7 million cubic meters per year, respectively, indicating that streamflow in the downstream portion of the basin has decreased more rapidly over time. Potential future research questions include the following: How do changes in precipitation compare with the expansion of irrigated areas in influencing streamflow within the river basin? How does the spatial distribution of expanded irrigated areas affect streamflow responses in different parts of the basin?

Other potential future research questions related to water withdrawal and river management policies are the following: How do different forms of water policies and human interventions affect long-term water availability, streamflow, and environmental health? Do the policies implemented by governments and water managers help mitigate indirect impacts, such as climate change? How do local water policies (such as




Figure 3. Downstream (solid line) and upstream (dashed line) annual streamflow for the Kissimmee River in the United States (Label B in Fig. 1). The basin’s river management is shown in different colors. For more information on this data and river, see Chen *et al.* (2016).




Figure 4. Downstream (solid line) and upstream (dashed line) streamflow in the Tarim River in China (Label C in Fig. 1). Irrigation and water reallocation policies are shown in color. For more information on the data and river, see Liu *et al.* (2014).

those in the Kissimmee and Tarim river basins) compare with regional policies (such as those in California and the Russian River)?

CHHAI can also be used to study the concept of anthropogenic drought (AghaKouchak *et al.* 2021), defined as drought events caused or intensified by human activities (Ashraf *et al.* 2021). The Tarim River basin (Fig. 4) is an example of anthropogenic drought, where downstream streamflow decreased even though precipitation increased in the basin (Yaning *et al.* 2009). The Helmand River basin in Afghanistan, Iran, and Pakistan (label “D” in Fig. 1) is another case of anthropogenic drought. The Hamun Lakes and Chah-nimehs are reservoirs located in the basin along the Iran–Afghanistan border. These lakes experienced a large increase in surface water storage in the early 1990s before experiencing

a substantial decrease in the late 1990s. During this time frame, the amount of land used for agriculture increased (Fig. 5). After the decrease in surface water storage, agricultural area continued to expand, but there was no clear corresponding change in surface water storage. Although the significant decrease in surface water storage coincided with a drought, the drought did not cause the decrease but instead accelerated it (Modaresi Rad *et al.* 2022). Future research questions for this basin include the following: What specific anthropogenic stressors contributed to the large decrease in storage around 2000? Why did surface water storage increase dramatically in the early 1990s?

Anthropogenic drought due to overuse of water resources has also caused saline lakes worldwide to shrink. One example of lake drought (Cheng *et al.* 2025) is Lake Urmia in Iran (label




Figure 5. Surface water storage of the Hamun Lakes and Chah-nimeh lakes and the area of land for agricultural use in the Helmand River basin in Afghanistan and Iran (Label D in Fig. 1). Dashed lines are the monthly data for the surface water storage (blue) and agricultural area (red) while solid lines are the 12-month rolling average. For further information on this data, see Modaresi Rad *et al.* (2022).




Figure 6. Surface water withdrawal due to agriculture in the basin (red) and lake water level above sea level (blue) for Urmia Lake in Iran (Label E in Fig. 1). For more information about the lake and data, see Alborzi *et al.* (2018).

“E” in Fig. 1; Alborzi *et al.* 2018; Pouladi *et al.* 2022). In the Lake Urmia basin, surface water withdrawal due to agriculture increased dramatically in the 2000s, and during the same time, the water level of the lake decreased consistently (Fig. 6). This increase and subsequent decrease in water level coincided with a similar pattern in the Helmand River basin (Fig. 5). As the basins are in the same region, there may be similar factors contributing to the decline in lake water levels. Further analysis of anthropogenic stressors in the region may reveal why the water storage and levels in the two basins declined so drastically.

Potential future research questions related to anthropogenic drought include the following: How do different types of human activities (e.g. increases in water demand) influence the onset, duration, and severity of local hydrologic droughts (e.g. see Pouladi *et al.* 2025b)? Can CHHAI help disentangle natural and anthropogenic drivers of recent historical droughts? Under what conditions do human interventions transform a meteorological drought (i.e. a deficit in precipitation) into a more severe hydrologic drought (i.e. a change in surface water storage; see Pouladi *et al.* 2025a)? What is the expected compounding impact of a meteorological drought and increasing water demand (e.g. see Mehran *et al.* 2017)?

In this section, we presented five case studies revolving around changing water policies and anthropogenic drought, and we outlined ideas for future research using CHHAI datasets related to those regions and topics. However, there are 20 other regions included in CHHAI, and we encourage researchers to analyze other human impacts, such as flood protection, land use changes, and reservoir implementation (see column 3 of Table A1). The hydrologic applications are not limited to the examples presented here. The dataset also supports comparative analyses across basins and time periods, enabling researchers to test hypotheses about how different management actions and socioeconomic pressures shape hydrologic

responses under similar climate conditions. Additionally, CHHAI can be used to identify patterns across different regions and basins. For example, similarities were observed in the timing of water level changes across Lake Urmia, Hamun Lakes, and the Chah-nimehs, and it is likely that these sorts of patterns could be discovered using the data provided by CHHAI.

Many basins included in CHHAI have been the subject of earlier studies, providing an opportunity to connect the benchmark dataset with established process understanding. For example, the Helmand River and Lake Urmia basins reflect well-documented cases of anthropogenic drought and maladaptation, driven by feedbacks among infrastructure development, governance, and human water use behavior (AghaKouchak *et al.* 2021, Pouladi *et al.* 2022). In transboundary river systems, such as the Mekong River basin, prior analyses have shown how cooperation and conflict coevolve with hydrological variability, infrastructure development, and institutional change (Lu *et al.* 2021, Wei *et al.* 2022). By building on these earlier insights and identifying patterns across regions, researchers can revisit known sociohydrological phenomena and human impacts with CHHAI by using standardized, comparable data across basins, thereby facilitating synthesis and hypothesis testing beyond individual case studies.

Conclusion

The hydrology community has created standardized datasets (e.g. MOPEX, CAMELS) for modeling hydrological systems in nonregulated basins, and they have also developed hydrological models incorporating human impacts such as water use and demand. However, there is a lack of standardized datasets combining hydrologic and human variables, which is why we present a dataset that provides observations of coupled human–water systems in basins across the world. Currently, the CHHAI

dataset consists of data for 25 regions from all continents, excluding Antarctica. The data reflects various human impacts on hydrological dynamics, such as land use changes, irrigation and other water uses, water policies, reservoir management and flood protection. With standardized data from basins across the world, there is potential to explore patterns in the challenges of coupled human–water systems across basins.

Although CHHAI consists of a variety of data types in different locations, it can be improved by including more data from Africa, Europe, and South America, as well as data that provides information on transboundary river management (e.g. Shrestha *et al.* 2022), river channelization (e.g. Chen *et al.* 2016), water diversion to and from rivers (e.g. Garcia *et al.* 2024), and flood protection (e.g. Paprotny *et al.* 2018, Nakamura *et al.* 2024). Currently, CHHAI does not include any data on topics such as social memory and human settlement patterns (Albertini *et al.* 2020), which would be a valuable inclusion. For this reason, CHHAI is a living dataset, and we encourage scholars performing research on coupled human–water systems to submit data to CHHAI to collaborate and share knowledge with others. As we receive more data, we will continue updating the dataset in future versions.

Beyond its immediate analytical applications, CHHAI can be viewed as a data legacy of the IAHS Panta Rhei Scientific Decade and as a foundation for future sociohydrological and coupled human–water systems research. Recent synthesis and perspective studies emphasize the need to move from isolated case studies toward shared benchmarks, comparative analyses, and solution-oriented research that explicitly links human behavior, institutions, and hydrological processes (Kreibich *et al.* 2023, Arheimer *et al.* 2024). Moreover, emerging work on societal value transitions highlights the importance of capturing long-term changes in norms, preferences, and governance alongside hydrological change (Wei *et al.* 2023). Together, these perspectives position CHHAI as an enabling platform for advancing comparative, policy-relevant, and locally grounded coupled human–water systems research aimed at improving water security and sustainability under global change.

Acknowledgements

This effort is inspired by the second and third International Association of Hydrological Sciences (IAHS) Scientific Decades: Panta Rhei – Everything Flows (2013–2022) and Hydrology Engaging Local People IN one Global world [HELPING; 20232032]. We acknowledge individuals and organizations that provided data for regions, including the Royal Irrigation Department, Thailand, which provided data for the Chao Phraya River basin.

Disclosure statement

No potential conflict of interest was reported by the author(s).

Funding

This material is supported by the United States National Science Foundation Graduate Research Fellowship Program under Grant No. DGE2235784 and by the National Science Foundation Office of International Science and Engineering Grant No. 2114701 and NSF Grant No. 2332263.

ORCID

Azara Boschee  <http://orcid.org/0009-0008-7475-2228>
 Augustina Clara Alexander  <http://orcid.org/0000-0002-0155-1678>
 Erhu Du  <http://orcid.org/0000-0002-0086-2394>
 Jesse Duku  <http://orcid.org/0009-0008-6983-7681>
 Aldo Fiori  <http://orcid.org/0000-0002-6662-5738>
 Margaret Garcia  <http://orcid.org/0000-0002-2864-2377>
 Heidi Kreibich  <http://orcid.org/0000-0001-6274-3625>
 Pedro Medeiros  <http://orcid.org/0000-0002-4879-3148>
 Joel Nobert  <http://orcid.org/0000-0002-6537-9950>
 Afua Owusu  <http://orcid.org/0000-0001-6420-6663>
 Mahendran Roobavannan  <http://orcid.org/0000-0003-1808-9720>
 Lukas Schoppa  <http://orcid.org/0000-0002-3443-4603>
 Ashish Shrestha  <http://orcid.org/0000-0002-0106-0318>
 Jiale Wang  <http://orcid.org/0009-0006-6413-8289>
 Amir AghaKouchak  <http://orcid.org/0000-0003-4689-8357>

Data availability

The CHHAI dataset is openly available on HydroShare at <https://doi.org/10.4211/hs.e011b48fabd2465eb5135275d2f3a93e>.

References

- Addor, N., *et al.*, 2017. The CAMELS data set: catchment attributes and meteorology for largesample studies. *Hydrology and Earth System Sciences*, 21 (10), 5293–5313. doi:10.5194/hess2152932017
- Addor, N. *et al.*, 2020. Large-sample hydrology: recent progress, guidelines for new datasets and grand challenges. *Hydrological Sciences Journal*, 65 (5), 712–725.
- AghaKouchak, A., *et al.*, 2021. Anthropogenic drought: definition, challenges, and opportunities. *Reviews of Geophysics*, 59 (2), e2019RG000683. doi:10.1029/2019RG000683
- Albertini, C., *et al.*, 2020. sociohydrological modelling: the influence of reservoir management and societal responses on flood impacts. *Water*, 12 (5), 1384. doi:10.3390/w12051384
- Alborzi, A., *et al.*, 2018. Climateinformed environmental inflows to revive a drying lake facing meteorological and anthropogenic droughts. *Environmental Research Letters*, 13 (8), 084010. doi:10.1088/17489326/aad246
- Alexander, K., 2025. Two dams in Northern California could be razed under PG&E plan [online]. *San Francisco Chronicle*. Available from: <https://www.sfchronicle.com/climate/article/twodamsnortherncaliforniarazedpgeplan20786824.php> [Accessed 28 July 2025].
- AlvarezGarretton, C., *et al.*, 2018. The CAMELSCL dataset: catchment attributes and meteorology for large sample studies – chile dataset. *Hydrology and Earth System Sciences*, 22 (11), 5817–5846. doi:10.5194/hess2258172018
- Arheimer, B., *et al.*, 2020. Global catchment modelling using WorldWide HYPE (WWH), open data, and stepwise parameter estimation. *Hydrology and Earth System Sciences*, 24 (2), 535–559. doi:10.5194/hess245352020
- Arheimer, B., *et al.*, 2024. The IAHS science for solutions decade, with hydrology engaging local people IN one global world (HELPING). *Hydrological Sciences Journal*, 69 (11), 1417–1435. doi:10.1080/02626667.2024.2355202
- Arnold, J.G., *et al.*, 1998. Large area hydrologic modeling and assessment part I: model development. *JAWRA Journal of the American Water Resources Association*, 34 (1), 73–89. doi:10.1111/j.17521688.1998.tb05961.x
- Ashraf, S., Nazemi, A., and AghaKouchak, A., 2021. Anthropogenic drought dominates groundwater depletion in Iran. *Scientific Reports*, 11 (1). doi:10.1038/s4159802188522y
- Barendrecht, M.H., *et al.*, 2019. The value of empirical data for estimating the parameters of a sociohydrological flood risk model. *Water Resources Research*, 55 (2), 1312–1336. doi:10.1029/2018WR024128

- Bartosova, A., *et al.*, 2019. Future socioeconomic conditions may have a larger impact than climate change on nutrient loads to the Baltic Sea. *Ambio*, 48 (11), 1325–1336. doi:10.1007/s13280019012435
- Blair, P. and Buytaert, W., 2016. Sociohydrological modelling: a review asking ‘why, what and how? *Hydrology and Earth System Sciences*, 20 (1), 443–478. doi:10.5194/hess204432016
- Burek, P., *et al.*, 2020. Development of the community water model (CWatM v1.04) – a high-resolution hydrological model for global and regional assessment of integrated water resources management. *Geoscientific Model Development*, 13 (7), 3267–3298. doi:10.5194/gmd1332672020
- Casale, F., *et al.*, 2021. Exploring future vulnerabilities of subalpine Italian regulated lakes under different climate scenarios: bottom-up vs top-down and CMIP5 vs CMIP6. *Journal of Hydrology: Regional Studies*, 38, 100973.
- Ceola, S., Mård, J., and Di Baldassarre, G., 2023. Drought and human mobility in Africa. *Earth’s Future*, 11 (12). doi:10.1029/2023EF003510
- Chagas, V.B.P., *et al.*, 2020. CAMELSBR: hydrometeorological time series and landscape attributes for 897 catchments in Brazil. *Earth System Science Data*, 12 (3), 2075–2096. doi:10.5194/essd1220752020
- Chen, X., *et al.*, 2016. From channelization to restoration: sociohydrologic modeling with changing community preferences in the Kissimmee River Basin, Florida. *Water Resources Research*, 52 (2), 1227–1244. doi:10.1002/2015WR018194
- Cheng, X., *et al.*, 2025. Global assessment and hotspots of lake drought. *Communications Earth & Environment*, 6 (1), 308. doi:10.1038/s43247025022852
- Consortium of Universitys for the Advancement of Hydrologic Sciences Inc. (CUAHSI), 2025. *HydroShare* [online]. Available from: <https://www.hydroshare.org/> [Accessed 1 Aug 2025].
- Coxon, G., *et al.*, 2020. CAMELSGB: hydrometeorological time series and landscape attributes for 671 catchments in Great Britain. *Earth System Science Data*, 12 (4), 2459–2483. doi:10.5194/essd1224592020
- Dang, H., *et al.*, 2022. Hydrologic balance and inundation dynamics of Southeast Asia’s largest inland lake altered by hydropower dams in the Mekong River basin. *Science of the Total Environment*, 831, 154833. doi:10.1016/j.scitotenv.2022.154833
- Dang, T.D., Chowdhury, A.F.M.K., and Galelli, S., 2020. On the representation of water reservoir storage and operations in largescale hydrological models: implications on model parameterization and climate change impact assessments. *Hydrology and Earth System Sciences*, 24 (1), 397–416. doi:10.5194/hess243972020
- Denaro, S., *et al.*, 2017. Informing the operations of water reservoirs over multiple temporal scales by direct use of hydrometeorological data. *Advances in Water Resources*, 103, 51–63. doi:10.1016/j.advwatres.2017.02.012
- Di Baldassarre, G., *et al.*, 2013. Sociohydrology: conceptualising human-flood interactions. *Hydrology and Earth System Sciences*, 17 (8), 3295–3303. doi:10.5194/hess1732952013
- Di Baldassarre, G., *et al.*, 2019. Sociohydrology: scientific challenges in addressing the sustainable development goals. *Water Resources Research*, 55 (8), 6327–6355. doi:10.1029/2018WR023901
- Do Nascimento, T.V.M., *et al.*, 2024. EStreams: an integrated dataset and catalogue of streamflow, hydroclimatic and landscape variables for Europe. *Scientific Data*, 11 (1), 879. doi:10.1038/s41597024037061
- Du, E., *et al.*, 2022. Evaluating distributed policies for conjunctive surface water-groundwater management in large river basins: water uses versus hydrological impacts. *Water Resources Research*, 58 (1). doi:10.1029/2021WR031352
- Duan, Q., *et al.*, 2006. Model parameter estimation experiment (MOPEX): an overview of science strategy and major results from the second and third workshops. *Journal of Hydrology*, 320 (1–2), 3–17. doi:10.1016/j.jhydrol.2005.07.031
- Fiener, P., Auerswald, K., and Van Oost, K., 2011. Spatiotemporal patterns in land use and management affecting surface runoff response of agricultural catchments—a review. *EarthScience Reviews*, 106 (1–2), 92–104. doi:10.1016/j.earscirev.2011.01.004
- Fiori, A., *et al.*, 2025. Relevance of the net natural inflow for investigating longterm hydrological changes in a managed lake catchment. *Journal of Hydrology*, 662, 133925. doi:10.1016/j.jhydrol.2025.133925
- Garcia, M., Mohajer Iravanloo, B., and Sivapalan, M., 2024. A diagnostic approach to modeling watersheds with human interference. *Journal of Hydrology*, 641, 131823. doi:10.1016/j.jhydrol.2024.131823
- Giuliani, M., *et al.*, 2019. Detecting the state of the climate system via artificial intelligence to improve seasonal forecasts and inform reservoir operations. *Water Resources Research*, 55 (11), 9133–9147. doi:10.1029/2019WR025035
- Giuliani, M., *et al.*, 2020. Datadriven control of water reservoirs using an emulator of the climate system. *IFAC PapersOnLine*, 53 (2), 16531–16536. doi:10.1016/j.ifacol.2020.12.771
- Giuliani, M. and Castelletti, A., 2024. Integrating inverse reinforcement learning and direct policy search for modeling multipurpose water reservoir systems. *IEEE Control Systems Letters*, 8, 1529–1534. doi:10.1109/LCSYS.2024.3411831
- Grogan, D.S., *et al.*, 2022. Water balance model (WBM) v.1.0.0: a scalable gridded global hydrologic model with watertracking functionality. *Geoscientific Model Development*, 15 (19), 7287–7323. doi:10.5194/gmd1572872022
- Hanasaki, N., *et al.*, 2018. A global hydrological simulation to specify the sources of water used by humans. *Hydrology and Earth System Sciences*, 22 (1), 789–817. doi:10.5194/hess227892018
- Horsburgh, J.S., *et al.*, 2016. HydroShare: sharing diverse environmental data types and models as social objects with application to the hydrology domain. *JAWRA Journal of the American Water Resources Association*, 52 (4), 873–889. doi:10.1111/17521688.12363
- Huang, J., *et al.*, 2025. Solving the robustness puzzle: the joint impact of optimization approach, robustness metrics, and scenarios on water resources management under deep uncertainty. *Journal of Environmental Management*, 373, 123540. doi:10.1016/j.jenvman.2024.123540
- Kandasamy, J., *et al.*, 2014. Sociohydrologic drivers of the pendulum swing between agricultural development and environmental health: a case study from Murrumbidgee River basin, Australia. *Hydrology and Earth System Sciences*, 18 (3), 1027–1041. doi:10.5194/hess1810272014
- Kim, Y., *et al.*, 2018. Assessment of regional threats to human water security adopting the global framework: a case study in South Korea. *Science of the Total Environment*, 637–638, 1413–1422. doi:10.1016/j.scitotenv.2018.04.420
- Konar, M., *et al.*, 2019. Expanding the scope and foundation of sociohydrology as the science of coupled human-water systems. *Water Resources Research*, 55 (2), 874–887. doi:10.1029/2018WR024088
- Kratzert, F., *et al.*, 2023. Caravan A global community dataset for large-sample hydrology. *Scientific Data*, 10 (1). doi:10.1038/s4159702301975w
- Kreibich, H., *et al.*, 2022. The challenge of unprecedented floods and droughts in risk management. *Nature*, 608 (7921), 80–86. doi:10.1038/s41586022049175
- Kreibich, H., *et al.*, 2023. Panta Rhei benchmark dataset: sociohydrological data of paired events of floods and droughts. *Earth System Science Data*, 15 (5), 2009–2023. doi:10.5194/essd1520092023
- Kreibich, H., *et al.*, 2025. Panta Rhei: a decade of progress in research on change in hydrology and society. *Hydrological Sciences Journal*, 70 (7), 1210–1236. doi:10.1080/02626667.2025.2469762
- Kummu, M., *et al.*, 2011. How close do we live to water? A global analysis of population distance to freshwater bodies. *PLoS ONE*, 6 (6), e20578. doi:10.1371/journal.pone.0020578
- Li, C., *et al.*, 2024. Highresolution land surface modeling of the irrigation effects on evapotranspiration over the Yellow River basin. *Journal of Hydrology*, 633, 130986. doi:10.1016/j.jhydrol.2024.130986
- Li, S., *et al.*, 2017a. Evolution of the vegetation system in the Heihe River basin in the last 2000 years. *Hydrology and Earth System Sciences*, 21 (8), 4233–4244. doi:10.5194/hess2142332017
- Li, X., *et al.*, 2021. Novel hybrid coupling of ecohydrology and socioeconomy at river basin scale: a watershed system model for the Heihe River basin. *Environmental Modelling & Software*, 141, 105058. doi:10.1016/j.envsoft.2021.105058
- Li, Y., Giuliani, M., and Castelletti, A., 2017b. A coupled human–natural system to assess the operational value of weather and climate services for agriculture. *Hydrology and Earth System Sciences*, 21 (9), 4693–4709. doi:10.5194/hess2146932017

- Liang, X., *et al.*, 1994. A simple hydrologically based model of land surface water and energy fluxes for general circulation models. *Journal of Geophysical Research: Atmospheres*, 99 (D7), 14415–14428. doi:10.1029/94JD00483
- Liu, D., *et al.*, 2015. A conceptual sociohydrological model of the coevolution of humans and water: case study of the Tarim River basin, western China. *Hydrology and Earth System Sciences*, 19 (2), 1035–1054. doi:10.5194/hess191035-2015
- Liu, M., *et al.*, 2008. Effects of land-use and land-cover change on evapotranspiration and water yield in China during 1900–2000. *JAWRA Journal of the American Water Resources Association*, 44 (5), 1193–1207. doi:10.1111/j.1752-1688.2008.00243.x
- Liu, Y., *et al.*, 2014. Socio-hydrologic perspectives of the co-evolution of humans and water in the Tarim River basin, Western China: the Taiji-Tire model. *Hydrology and Earth System Sciences*, 18 (4), 1289–1303. doi:10.5194/hess-18-1289-2014
- Lu, Z., *et al.*, 2015a. Trade-offs between midstream agricultural production and downstream ecological sustainability in the Heihe River basin in the past half century. *Agricultural Water Management*, 152, 233–242. doi:10.1016/j.agwat.2015.01.022
- Lu, Z., *et al.*, 2015b. Evolution of the human–water relationships in the Heihe River basin in the past 2000 years. *Hydrology and Earth System Sciences*, 19 (5), 2261–2273. doi:10.5194/hess-19-2261-2015
- Lu, Z., *et al.*, 2018. Co-evolutionary dynamics of the human–environment system in the Heihe River basin in the past 2000 years. *Science of the Total Environment*, 635, 412–422. doi:10.1016/j.scitotenv.2018.03.231
- Lu, Y., *et al.*, 2021. Socio-hydrologic modeling of the dynamics of cooperation in the transboundary Lancang–Mekong River. *Hydrology and Earth System Sciences*, 25 (4), 1883–1903. doi:10.5194/hess-25-1883-2021
- Mai, J., *et al.*, 2022. The Great Lakes runoff intercomparison project phase 4: the Great Lakes (GRIP-GL). *Hydrology and Earth System Sciences*, 26 (13), 3537–3572. doi:10.5194/hess-26-3537-2022
- Maity, R., *et al.*, 2024. Revolutionizing the future of hydrological science: impact of machine learning and deep learning amidst emerging explainable AI and transfer learning. *Applied Computing and Geosciences*, 24, 100206. doi:10.1016/j.acags.2024.100206
- Medeiros, P. and Sivapalan, M., 2020. From hard-path to soft-path solutions: slow–fast dynamics of human adaptation to droughts in a water scarce environment. *Hydrological Sciences Journal*, 65 (11), 1803–1814. doi:10.1080/02626667.2020.1770258
- Mehran, A., *et al.*, 2017. Compounding impacts of human-induced water stress and climate change on water availability. *Scientific Reports*, 7 (1), 6282. doi:10.1038/s41598-017-06765-0
- Meira Neto, A.A., *et al.*, 2024. Evolution of drought mitigation and water security through 100 years of reservoir expansion in semi-arid Brazil. *Water Resources Research*, 60 (9), e2023WR036411. doi:10.1029/2023WR036411
- Modaresi Rad, A., *et al.*, 2022. Anthropogenic stressors compound climate impacts on inland lake dynamics: the case of Hamun Lakes. *Science of the Total Environment*, 829, 154419. doi:10.1016/j.scitotenv.2022.154419
- Montanari, A., *et al.*, 2013. “Panta Rhei—everything flows”: change in hydrology and society—the IAHS scientific decade 2013–2022. *Hydrological Sciences Journal*, 58 (6), 1256–1275. doi:10.1080/02626667.2013.809088
- Mostert, E., 2018. An alternative approach for socio-hydrology: case study research. *Hydrology and Earth System Sciences*, 22 (1), 317–329. doi:10.5194/hess-22-317-2018
- Müller Schmied, H., *et al.*, 2021. The global water resources and use model WaterGAP v2.2d: model description and evaluation. *Geoscientific Model Development*, 14 (2), 1037–1079. doi:10.5194/gmd-14-1037-2021
- Nakamura, S., *et al.*, 2024. Levee system transformation in coevolution between humans and water systems along the Kiso River, Japan. *Hydrology and Earth System Sciences*, 28 (10), 2329–2342. doi:10.5194/hess-28-2329-2024
- Newman, A.J., *et al.*, 2015. Development of a large-sample watershed-scale hydrometeorological data set for the contiguous USA: data set characteristics and assessment of regional variability in hydrologic model performance. *Hydrology and Earth System Sciences*, 19 (1), 209–223. doi:10.5194/hess-19-209-2015
- Odongo, V.O., *et al.*, 2019. Impact of land use and land cover transitions and climate on evapotranspiration in the Lake Naivasha Basin, Kenya. *Science of the Total Environment*, 682, 19–30. doi:10.1016/j.scitotenv.2019.04.062
- Oki, T. and Kanae, S., 2006. Global hydrological cycles and world water resources. *Science*, 313 (5790), 1068–1072. doi:10.1126/science.1128845
- Owusu, A., *et al.*, 2025. Bridging scales and borders on water availability and use in the transboundary Volta River Basin: a water accounting approach. *Journal of Hydrology: Regional Studies*, 59, 102377.
- Padiyedath Gopalan, S., *et al.*, 2021. Impact assessment of reservoir operation in the context of climate change adaptation in the Chao Phraya River basin. *Hydrological Processes*, 35 (1), e14005. doi:10.1002/hyp.14005
- Padiyedath Gopalan, S., *et al.*, 2022a. Inclusion of flood diversion canal operation in the H08 hydrological model with a case study from the Chao Phraya River basin: model development and validation. *Hydrology and Earth System Sciences*, 26 (9), 2541–2560. doi:10.5194/hess-26-2541-2022
- Padiyedath Gopalan, S., *et al.*, 2022b. Potential impact of diversion canals and retention areas as climate change adaptation measures on flood risk reduction: a hydrological modelling case study from the Chao Phraya River Basin, Thailand. *Science of the Total Environment*, 841, 156742. doi:10.1016/j.scitotenv.2022.156742
- Padiyedath Gopalan, S., *et al.*, 2024. Adaptation portfolio – a multi-measure framework for future floods and droughts. *Npj Natural Hazards*, 1 (1), 48. doi:10.1038/s44304-024-00048-1
- Pan, H., *et al.*, 2018. Sociohydrology modeling for complex urban environments in support of integrated land and water resource management practices. *Land Degradation & Development*, 29 (10), 3639–3652. doi:10.1002/ldr.3106
- Pande, S. and Sivapalan, M., 2017. Progress in socio-hydrology: a meta-analysis of challenges and opportunities. *WIREs Water*, 4 (4). doi:10.1002/wat2.1193
- Paprotny, D., Morales-Nápoles, O., and Jonkman, S.N., 2018. HANZE: a pan-European database of exposure to natural hazards and damaging historical floods since 1870. *Earth System Science Data*, 10 (1), 565–581. doi:10.5194/essd-10-565-2018
- Peel, M.C. and Blöschl, G., 2011. Hydrological modelling in a changing world. Progress in physical geography. *Earth and Environment*, 35 (2), 249–261.
- Pereira, B., *et al.*, 2025a. Advancing water security in semi-arid Brazil: expansion of water storage infrastructure and human–water system coevolution. *Frontiers in Water*, 7, 1602146. doi:10.3389/frwa.2025.1602146
- Pereira, B.S., *et al.*, 2025b. Hydrological heritage: a historical exploration of human–water dynamics in northeast Brazil. *Hydrological Sciences Journal*, 70 (4), 546–558. doi:10.1080/02626667.2024.2446272
- Pouladi, M., *et al.*, 2025a. Addressing data limitations to explore management strategies and adaptations using stylized agent-based modeling: a case study of socio-environmental systems. *Ecological Modelling*, 501, 110997. doi:10.1016/j.ecolmodel.2024.110997
- Pouladi, P., *et al.*, 2022. Desiccation of a saline lake as a lock-in phenomenon: a socio-hydrological perspective. *Science of the Total Environment*, 811, 152347. doi:10.1016/j.scitotenv.2021.152347
- Pouladi, P., *et al.*, 2025b. Socio-hydrological lock-in; an emergent phenomenon in the face of anthropogenic drought. *Journal of Environmental Management*, 373, 123318. doi:10.1016/j.jenvman.2024.123318
- Reed, P.M., *et al.*, 2022. Multisector dynamics: advancing the science of complex adaptive human–earth systems. *Earth’s Future*, 10 (3), 3. doi:10.1029/2021EF002621
- Reed, S., *et al.*, 2004. Overall distributed model intercomparison project results. *Journal of Hydrology*, 298 (1–4), 27–60. doi:10.1016/j.jhydrol.2004.03.031
- Ricart, S., Gandolfi, C., and Castelletti, A., 2024. Targeting farmers’ heterogeneity to enrich climate change adaptation policy design: findings from northern Italy. *Environmental Research: Climate*, 3 (3), 031001.

- Ricart, S., Gandolfi, C., and Castelletti, A., 2025. What drives farmers' behavior under climate change? Decoding risk awareness, perceived impacts, and adaptive capacity in northern Italy. *Heliyon*, 11 (1), e41328. doi:10.1016/j.heliyon.2024.e41328
- Rose, S. and Peters, N.E., 2001. Effects of urbanization on streamflow in the Atlanta area (Georgia, USA): a comparative hydrological approach. *Hydrological Processes*, 15 (8), 1441–1457. doi:10.1002/hyp.218
- Salvadore, E., Bronders, J., and Batelaan, O., 2015. Hydrological modeling of urbanized catchments: a review and future directions. *Journal of Hydrology*, 529, 62–81. doi:10.1016/j.jhydrol.2015.06.028
- Schaake, J., Cong, S., and Duan, Q., 2006. U.S. MOPEX DATA SET. *IAHS Publication Series*, 307 (N/A), 9–28.
- Shrestha, A., et al., 2022. Socio-hydrological modeling of the tradeoff between flood control and hydropower provided by the Columbia River Treaty. *Hydrology and Earth System Sciences*, 26 (19), 4893–4917. doi:10.5194/hess-26-4893-2022
- Sieber, J., 2006. WEAP water evaluation and planning system. Presented at the 3rd International Congress on Environmental Modelling and Software, Burlington, Vermont, USA.
- Sivapalan, M. and Blöschl, G., 2015. Time scale interactions and the coevolution of humans and water: time scale interactions and coevolution of humans and water. *Water Resources Research*, 51 (9), 6988–7022. doi:10.1002/2015WR017896
- Sivapalan, M., Savenije, H.H.G., and Blöschl, G., 2012. Socio-hydrology: a new science of people and water. *Hydrological Processes*, 26 (8), 1270–1276. doi:10.1002/hyp.8426
- Smith, M., et al., 2013. The distributed model intercomparison project – phase 2: experiment design and summary results of the western basin experiments. *Journal of Hydrology*, 507, 300–329. doi:10.1016/j.jhydrol.2013.08.040
- Sousa, D.S., et al., 2025. Coupling data-driven agent-based and hydrological modelling to explore the effect of collective water allocation strategies in environmental flows. *Journal of Hydrology*, 652, 132670. doi:10.1016/j.jhydrol.2025.132670
- Sui, X. and Van de Ven, F.H.M., 2023. The influence of low impact development (LID) on basin runoff in a half-urbanized catchment: a case study in San Antonio, Texas. *Journal of Hydrology*, 616, 128793. doi:10.1016/j.jhydrol.2022.128793
- Sunkara, S.V., Singh, R., and Bhave, A., 2025. Quantifying the value of stakeholder-elicited information in models of coupled human–water systems. *Hydrological Sciences Journal*, 70 (3), 467–482. doi:10.1080/02626667.2024.2437131
- Sutanudjaja, E.H., et al., 2018. PCR-GLOBWB 2: a 5 arcmin global hydrological and water resources model. *Geoscientific Model Development*, 11 (6), 2429–2453. doi:10.5194/gmd-11-2429-2018
- Tian, F., et al., Eds., 2026. *Coevolution and prediction of coupled human-water systems*. Amsterdam, Netherlands: Elsevier.
- Tijerina-Kreuzer, D., et al., 2021. Continental hydrologic intercomparison project, phase I: a large-scale hydrologic model comparison over the continental United States. *Water Resources Research*, 57 (7). doi:10.1029/2020WR028931
- Trabelsi, F. and Bel Hadj Ali, S., 2022. Exploring machine learning models in predicting irrigation groundwater quality indices for effective decision making in Medjerda River Basin, Tunisia. *Sustainability*, 14 (4), 2341. doi:10.3390/su14042341
- Trabelsi, F., Bel Hadj Ali, S., and Lee, S., 2023. Comparison of novel hybrid and benchmark machine learning algorithms to predict groundwater potentiality: case of a Drought-Prone Region of Medjerda Basin, Northern Tunisia. *Remote Sensing*, 15 (1), 152. doi:10.3390/rs15010152
- Troy, T.J., et al., 2015a. Moving sociohydrology forward: a synthesis across studies. *Hydrology and Earth System Sciences*, 19 (8), 3667–3679. doi:10.5194/hess-19-3667-2015
- Troy, T.J., Pavao-Zuckerman, M., and Evans, T.P., 2015b. Debates—Perspectives on socio-hydrology: socio-hydrologic modeling: tradeoffs, hypothesis testing, and validation. *Water Resources Research*, 51 (6), 4806–4814. doi:10.1002/2015WR017046
- Turner, S., et al., 2025. ROBIN: reference observatory of basins for international hydrological climate change detection. *Scientific Data*, 12 (1), 654. doi:10.1038/s41597-025-04907-y
- van Beek, L.P.H. and Bierkens, M.F.P., 2008. *The global hydrological model PCR-GLOBWB: conceptualization, parameterization and verification*. The Netherlands: Department of Physical Geography, Utrecht University.
- Wada, Y., et al., 2017. Human–water interface in hydrological modelling: current status and future directions. *Hydrology and Earth System Sciences*, 21 (8), 4169–4193. doi:10.5194/hess-21-4169-2017
- Wei, Y., et al., 2022. A socio-hydrological framework for understanding conflict and cooperation with respect to transboundary rivers. *Hydrology and Earth System Sciences*, 26 (8), 2131–2146. doi:10.5194/hess-26-2131-2022
- Wei, J., et al., 2023. Transition in the societal value and governance of water resources in Australia and China. *Humanities and Social Sciences Communications*, 10 (1), 359. doi:10.1057/s41599-023-01857-x
- Wisser, D., et al., 2008. Global irrigation water demand: variability and uncertainties arising from agricultural and climate data sets. *Geophysical Research Letters*, 35 (24). doi:10.1029/2008GL035296
- Wu, J., et al., 2021. Reservoirs regulate the relationship between hydrological drought recovery water and drought characteristics. *Journal of Hydrology*, 603, 127127. doi:10.1016/j.jhydrol.2021.127127
- Yang, G., Giuliani, M., and Castelletti, A., 2023. Operationalizing equity in multipurpose water systems. *Hydrology and Earth System Sciences*, 27 (1), 69–81. doi:10.5194/hess-27-69-2023
- Yanling, C., et al., 2009. Fifty-year climate change and its effect on annual runoff in the Tarim River Basin, China. *Quaternary International*, 208 (1–2), 53–61. doi:10.1016/j.quaint.2008.11.011
- Ye, Z., Chen, Y., and Zhang, X., 2014. Dynamics of runoff, river sediments and climate change in the upper reaches of the Tarim River, China. *Quaternary International*, 336, 13–19. doi:10.1016/j.quaint.2013.06.006
- Zhang, Y., et al., 2021. Coupled hydrology-crop growth model incorporating an improved evapotranspiration module. *Agricultural Water Management*, 246, 106691. doi:10.1016/j.agwat.2020.106691
- Zhao, Y., et al., 2016. Downstream ecosystem responses to middle reach regulation of river discharge in the Heihe River Basin, China. *Hydrology and Earth System Sciences*, 20 (11), 4469–4481. doi:10.5194/hess-20-4469-2016

Appendix A. Information about the data

Table A1. Overview of CHHA1 data from 25 study regions. The third column indicates the type of variables that are in the dataset. For example, “Meteorological” variables may include precipitation, evapotranspiration, temperature, and other meteorological variables. The fifth column shows the name of the folder in the HydroShare Resource where the basin’s data can be found.

Basin/study region	Country	Types of variables	Data types	Time period covered	Dataset ZIP folder	Citation(s)
Helmand River	Afghanistan	Reservoir, meteorological, land use	Monthly time series	1984–2019	AFG_basin_HelmandRiver	Modaresi Rad <i>et al.</i> 2022
Murrumbidge River	Australia	Reservoir, meteorological, socioeconomic, crop production	Daily time series, yearly time series	1967–2012	AUS_basin_MurrumbidgeeRiver	Kandasamy <i>et al.</i> 2014
Formoso River	Brazil	Streamflow, other hydrological, water use, irrigation, cooperation toward water withdrawal rules	Daily time series, shapefiles	2016–2022	BRA_basin_FormosoRiver	Sousa <i>et al.</i> 2025
Jaguaribe River	Brazil	Streamflow, reservoir, meteorological, land use, water use	Yearly time series, Shapefiles	1872–2022	BRA_basin_JaguaribeRiver	Medeiros and Sivapalan 2020, Meira Neto <i>et al.</i> 2024, Pereira <i>et al.</i> 2025a, 2025b
Columbia River	Canada, United States	Streamflow, reservoir, transboundary cooperation	Monthly time series, daily time series	1990–2017	CAN_basin_ColumbiaRiver	Shrestha <i>et al.</i> 2022
Dongjiang River	China	Reservoir, streamflow	Monthly time series	1956–2011	CHN_basin_DongjiangRiver	Wu <i>et al.</i> 2021
Heihe River	China	Streamflow, meteorological, land use, water use, crop production	Daily time series, yearly time series, historical	1958–2016	CHN_basin_HeiheRiver	Lu <i>et al.</i> 2015a, 2015b, Zhao <i>et al.</i> 2016, Li <i>et al.</i> 2017a, 2021, Lu <i>et al.</i> 2018, Du <i>et al.</i> 2022
Tarim River	China	Meteorological, streamflow, land use, socioeconomic, river management	Yearly time series	1937–2025	CHN_basin_TarimRiver	Liu <i>et al.</i> 2014, 2015
Yellow River	China	Water use, water diversion	Yearly time series	2003–2020	CHN_basin_YellowRiver	Li <i>et al.</i> 2024
Elbe River	Germany and Czech Republic	Streamflow, meteorological, socioeconomic, land use, flood protection	Daily time series, yearly time series, dam information	1978–2020	DEU_basin_ElbeRiver	Paprotny <i>et al.</i> 2018, Barendrecht <i>et al.</i> 2019
Volta River	Ghana	Streamflow, meteorological, land use	Yearly time series	2003–2021	GHA_basin_VoltaRiver	Owusu <i>et al.</i> 2025
Krishna River	India	Groundwater, reservoir, meteorological, water demand	Monthly time series	1968–2013	IND_basin_KrishnaRiver	Sunkara <i>et al.</i> 2025
Lake Urmia	Iran	Other hydrological, meteorological, population, economic, land use, water use	Monthly time series, yearly time series	1955–2020	IRN_basin_LakeUrmia	Alborzi <i>et al.</i> 2018, Pouladi <i>et al.</i> 2022
Lake Bracciano	Italy	Streamflow, water use, meteorological, other hydrological	Yearly time series, daily time series	1921–2021	ITA_basin_LakeBracciano	Fiori <i>et al.</i> 2025

(Continued)

Table A1. (Continued).

Basin/study region	Country	Types of variables	Data types	Time period covered	Dataset ZIP folder	Citation(s)
Lake Como	Italy	Reservoir, economic, meteorological, irrigation, other hydrological	Daily time series, hourly time series, sub-hourly time series	2006–2022	ITA_basin_LakeComo	Denaro <i>et al.</i> 2017, Giuliani <i>et al.</i> 2019, Giuliani <i>et al.</i> 2020, Casale <i>et al.</i> 2021, Yang <i>et al.</i> 2023, Giuliani and Castelletti 2024, Ricart <i>et al.</i> 2024, Huang <i>et al.</i> 2025, Ricart <i>et al.</i> 2025
Kiso River Basin	Japan	Flood protection, land use	Shapefiles	1890–2009	JPN_basin_KisoRiver	Nakamura <i>et al.</i> 2024
Republic of Korea	Republic of Korea	Land use, water use	NetCDF, yearly time series, spatial data	2000–2010	KOR_region_SouthKorea	Kim <i>et al.</i> 2018
Chao Phraya River	Thailand	Reservoir, canals, streamflow, water use	Daily time series, monthly time series	1993–2019	THA_basin_ChaoPhrayaRiver	Padiyedath Gopalan <i>et al.</i> 2021, 2022a, 2022b, 2024
Medjerda River	Tunisia	Precipitation, land use	Daily time series, yearly time series, shapefiles, GeoTIFFs	1905–2020	TUN_basin_MedjerdaRiver	Trabelsi and Bel Hadj Ali 2022, Trabelsi <i>et al.</i> 2023
Rufiji River	Tanzania	Water use	Information by location	Static in time	TZA_basin_RufijiRiver	–
Lake Lanier	United States	Reservoir	Daily time series	1948–2012	USA_basin_LakeLanier	–
Russian River	United States	Streamflow, irrigation, meteorological, river management	Daily time series	1959–2022	USA_basin_RussianRiver	Garcia <i>et al.</i> 2024
Kissimmee River	United States	Streamflow, meteorology, population, economic, river management	Daily time series, yearly time series	1942–2013	USA_basin_KissimmeeRiver	Chen <i>et al.</i> 2016
Sacramento River	United States	Reservoir	Daily time series	2000–2023	USA_basin_SacramentoRiver	–
Mekong River	Vietnam, Cambodia, Thailand, Laos, Myanmar	Crop, meteorological, irrigation, land use, economic, population, streamflow, other hydrological	Shapefiles, netCDFs, GeoTIFFs, dam information, other gridded data	1961–2022	VNM_basin_MekongRiver	Dang <i>et al.</i> 2022